



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

Homocysteine, folate and vitamin B12 in neuropsychiatric diseases: review and treatment recommendations

Stanger, Olaf ; Fowler, Brian ; Pietrzik, Klaus ; Huemer, Martina ; Haschke-Becher, Elisabeth ;
Semmler, Alexander ; Lorenzl, Stefan ; Linnebank, Michael

Abstract: In Europe, neuropsychiatric diseases currently make up approximately a third of the total burden of disease. In 2004, 27% of the overall population was affected by at least one of the most frequent neuropsychiatric diseases such as Alzheimer's dementia, Parkinson's disease, stroke or depression. The annual costs of care exceed those of cancer, cardiovascular conditions and diabetes. In order to delay the onset or course of neurodegenerative diseases, the available potential should be utilized. As well as improving quality of life of patients and relatives, this may reduce the great financial burden caused by neurodegenerative disorders. However, the availability of established drugs or therapeutic agents is very limited. This paper reviews the state of current knowledge as to how homocysteine metabolism is relevant for neurodegenerative and other neuropsychiatric diseases, with particular emphasis on the evidence for prophylactic and therapeutic strategies. In the European countries, many people do not take the recommended daily minimum amount of folate and vitamin B12. Deficiency of these vitamins and secondary changes in the concentrations of associated metabolites, such as methylmalonic acid and homocysteine, may contribute to the onset and progression of neuropsychiatric diseases. This paper reviews the evidence regarding whether substitution of folate and vitamin B12 is beneficial, for example, in cerebrovascular disease, dementia and depression.

DOI: <https://doi.org/10.1586/ern.09.75>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-21178>

Journal Article

Accepted Version

Originally published at:

Stanger, Olaf; Fowler, Brian; Pietrzik, Klaus; Huemer, Martina; Haschke-Becher, Elisabeth; Semmler, Alexander; Lorenzl, Stefan; Linnebank, Michael (2009). Homocysteine, folate and vitamin B12 in neuropsychiatric diseases: review and treatment recommendations. *Expert Review of Neurotherapeutics*, 9(9):1393-1412.

DOI: <https://doi.org/10.1586/ern.09.75>

Homocysteine, folate and vitamin B12 in neuropsychiatric diseases: review of the literature and recommendations for treatment

*Olaf Stanger, Brian Fowler, Klaus Pietrzik, Martina Huemer, Elisabeth Haschke-Becher, Alexander Semmler, Stefan Bleich, Stefan Lorenzl, Michael Linnebank**

***corresponding author**

Olaf Stanger , Prof. Dr.	Paracelsus Medical University Salzburg, Clinic for Heart Surgery, ÖGK Institute for Applied Aminoacid and Vitamin Research, Müllner Hauptstrasse 48, A-5020 Salzburg. Email: o.stanger@salk.at
Michael Linnebank , Dr.	Zurich University, University Hospital, Clinic for Neurology, Frauenklinikstrasse 26, CH-8091 Zurich. Email: michael.linnebank@usz.ch
Brian Fowler , Prof. Dr.	Basle University Children's Hospital (UKBB), Metabolism Department. Römergasse 8, Postfach, CH-4005 Basle. Email: Brian.Fowler@ukbb.ch
Klaus Piertzik , Prof. Dr.	Rheinische Friedrich-Wilhelm University of Bonn, Institute for Nutrition and Food Sciences. Endenicher Allee 11-13, D-53115 Bonn. Email: k.pietrzik@uni-bonn.de

- 1 **Martina Huemer**, Dr. Landeskrankenhaus Hospital Bregenz, Department for
2 Paediatrics and Adolescent Medicine. Carl-Pedenz-Str.
3 2, A-6900 Bregenz
4 Email: martina.huemer@lkhb.at.
5
- 6 **Elisabeth Haschke-Becher**, Dr. Paracelsus Medical University Salzburg, University
7 Clinic for Neurology, Central Laboratory of the
8 Christian-Doppler Hospital Salzburg. Ignaz-Harrer-Str.
9 79, A-5020 Salzburg. Email:
10 elisabeth.haschke@elisabethinen.or.at
11
- 12 **Stefan Lorenzl**, Priv.-Doz., Dr. Interdisciplinary Centre for Palliative Medicine
13 and Clinic for Neurology, LMU Hospital, Campus
14 Grosshadern. Marchioninistrasse 15, D-81377
15 Munich. Email: [Stefan.Lorenzl@med.uni-](mailto:Stefan.Lorenzl@med.uni-muenchen.de)
16 [muenchen.de](mailto:Stefan.Lorenzl@med.uni-muenchen.de)
17
- 18 **Alexander Semmler**, Dr. Zurich University, University Hospital, Clinic for
19 Neurology (Neurolab), August-Forel-Strasse 7, 8008
20 Zurich. Email: alexander.semmler@usz.ch

Abstract

In Europe, neuropsychiatric diseases currently make up around one third of the total burden of disease. In 2004, 27% of the overall population was affected by at least one of the most frequent neuropsychiatric diseases such as Alzheimer dementia, Parkinson's disease, stroke or depression. The annual costs of care exceed those of cancer, cardiovascular conditions and diabetes. In order to delay onset or course of neurodegenerative diseases, the available potential should be utilized. As well as improving quality of life of patients and relatives, this may reduce the great financial burden caused by neurodegenerative disorders. However, the availability of established drugs or therapeutic agents is very limited. This paper reviews the state of current knowledge on to what extent homocysteine metabolism is relevant for neurodegenerative and other neuropsychiatric diseases with particular emphasis on the evidence for prophylactic and therapeutic strategies.

In the European countries, many people do not take the recommended daily minimum amount of folate and vitamin B12. Deficiency of these vitamins and secondary changes in the concentrations of associated metabolites, such as methylmalonic acid and homocysteine, may contribute to onset and progression of neuropsychiatric diseases. This paper reviews the evidence on whether substitution of folate and vitamin B12 is beneficial e.g. in several cases of cerebrovascular disease, dementia and depression.

Introduction

Worldwide, around 400 million people suffer from neurological and mental disorders [1]. The general consensus is that neuropsychiatric diseases make up around 35%

of the total burden of disease in Europe [2,3]. For example, in Europe, around 1.1 million new cases of cerebral ischaemia are reported every year [4]. The annual treatment costs for a total of 127 million Europeans suffering from at least one of the most frequent neuropsychiatric diseases amounted to €386 billion in 2004, which is more than the costs caused by cancer, cardiovascular disorders or diabetes [2]. Apart from those direct costs for hospitalisation, medical and nursing care, rehabilitation, drugs, psycho- and physiotherapy, laboratory costs and medical-technical services, there are additional indirect costs due to incapacity, disability and early death.

In European countries, many people do not take the recommended daily minimum amount of vitamin B12 and folate (naturally occurring biologically active form of the vitamin) or folic acid (precursor of folate, e.g. ingredient of synthetic vitamin drugs) [5-7]. Deficiency of these vitamins and consecutive changes in the concentration of associated metabolites such as methylmalonic acid (MMA) and homocysteine may contribute to the onset and progression of neuropsychiatric diseases [8-15].

1. Metabolism

1.1 Biochemistry

Folate and vitamin B12 play an important role in the development, differentiation and function of the central nervous system. Both vitamins are involved in methionine-homocysteine metabolism (figure 1). Methionine becomes activated to S-adenosylmethionine (SAM), which is indispensable for numerous reactions involving methylation, e.g. in the synthesis of nucleic acids [DNA, RNA], proteins, neurotransmitters, hormones, fatty acids, polysaccharides, phospholipids or DNA

1 methylation [16]. For example, several neuroendocrinologically important micro-
2 molecules, such as noradrenaline and N-acetylserotonine, are converted by SAM
3 dependent methylation reactions into biologically active neurotransmitters [17].
4 Others, such as L-DOPA, can be deactivated through SAM-dependent methylations.
5 S-adenosylhomocysteine (SAH) results from SAM through the release of the methyl
6 group. Due to antagonism, elevated SAH concentrations reduce the SAM-dependent
7 methylation capacity and, by doing so, impair numerous metabolic processes in the
8 brain [18]. Several cell dysfunctions, DNA damage and disturbed biosynthesis of
9 myelin are potential consequences [19]. SAH is hydrolyzed into homocysteine. The
10 reaction converting SAH into homocysteine is reversible and is favoured, when an
11 increase in homocysteine levels occurs. Plasma homocysteine correlates closely
12 with the SAH level in the cerebrospinal fluid and the brain in animals [20].
13 Homocysteine is a neuro- and vasculotoxic sulphur-containing intermediary product.
14 Homocysteine can be transsulfurated to cystathionine and, subsequently, to
15 cysteine, which is a component of glutathione. Transsulfuration of homocysteine
16 depends on vitamin B6. Alternatively, homocysteine can be remethylated to
17 methionine by addition of a methyl group from 5-methyltetrahydrofolate (5-MTHF),
18 which is synthesized by 5,10-methylenetetrahydrofolate reductase (MTHFR).
19 Remethylation can be catalyzed by methionine-synthase (MS), which requires
20 vitamin B12 in the form of methylcobalamin as cofactor. By the remethylation
21 reaction of homocysteine to methionine, 5-MTHF is regenerated to 5,10-MTHF,
22 which is necessary for nucleic acid synthesis. Thus, vitamin B12 deficiency leading
23 to a reduced methylation rate of homocysteine to methionine can lead to a functional
24 deficiency of 5,10-MTHF and subsequent dysfunction e.g. of the haematopoietic
25 system ("folate trap"). Reduced remethylation of homocysteine to methionine and

SAM due to a lack of vitamin B12 or folate can yield elevated levels of homocysteine. Reduced synthesis of SAM can lead to a state of “hypomethylation” e.g. resulting in disturbed synthesis of neurotransmitters and proteins important for the structural integrity of the brain (figure 1) [21,22].

The homocysteine metabolism in the brain is different from the systemic homocysteine metabolism. In the liver and the kidneys, there is an alternative pathway available for remethylation, apart from MS, the enzyme betaine-homocysteine-methyltransferase (BHMT), but this enzyme has not been detected in the brain [23]. Alternatively, in other organs such as liver, kidney and the gastrointestinal tract, homocysteine can be irreversibly broken down into cysteine and glutathione via cystathionine through condensation with serine (transsulfuration). The activities of the two involved enzymes cystathionine- β -synthase (CBS) and γ -cystathionase are each dependent on vitamin B6 as a cofactor. Similar to endothelial cells, neurons and other CNS cells do not seem to strongly express the complete homocysteine transsulfuration pathway, i.e., the activity of γ -cystathionase is low, although some brain transsulfuration activity has been reported, recently [24,25]. Hence, the capacity of homocysteine metabolism in the CNS is largely dependent on sufficient supplies of folate and vitamin B12 [26]. In particular, glia cells only have very small vitamin B12 stores that are quickly exhausted in cases of negative balance [27].

Due to the cofactor function of adenosylcobalamin for the mitochondrial methylmalonyl-CoA-mutase, vitamin B12 deficiency leads to the conversion of methylmalonyl-CoA to methylmalonic acid, which can be neurotoxic [8,28]. Another

CNS-specific feature is the dependence on the transport of folates through the blood-brain barrier. During this active transport process, 5-MTHF at the choroid plexus binds to folate receptor proteins and reaches the neurons through endocytosis, storage and release via the cerebrospinal fluid compartment [29,30]. If active folate transport or metabolism at the choroid plexus is disturbed, 5-MTHF levels can become low in the cerebrospinal fluid even in the presence of normal plasma folate concentrations [31]. By active transport, 5-MTHF occurs in higher concentrations in the cerebrospinal fluid [around 14-18 ng/mL] than in the blood [3-12 ng/mL] [32]. Similarly, there exist active transport mechanisms for vitamin B6 and B12 [33,34].

1.2 Genetics

In addition to the availability of vitamins, genetic variants contribute to the inter-individual differences in homocysteine metabolism [35,36]. Several of these variants have been reported to be associated with neuropsychiatric diseases. However, literature is conflicting, and numerous studies did not observe any association of genetic variants of homocysteine metabolism with disease. The frequent missense variant of methylenetetrahydrofolate reductase (MTHFR) c.677C>T is (A222V) is associated with reduced enzyme activity, and homozygous carriers of this variant have a mean increase of plasma homocysteine levels of approximately 25%, whereas the effect on homocysteine levels is generally stronger when folate plasma levels are low [37-42]. Despite the association between MTHFR c.677TT and elevated homocysteine plasma levels, the association of this variant with different neuropsychiatric diseases is inconsistently reported in numerous studies [43-45]. Because homocysteine levels are modulated by factors like age, gender, smoking

1 and renal function, the homocysteine increasing effect of MTHFR c.677C>T might be
2 overridden or confounded by such factors [46-49]. Nevertheless, large meta-
3 analyses have proven that the T-allele is associated with cardio- and
4 cerebrovascular disease, and this polymorphism may also be associated with the
5 incidence of dementia [50-53].

6
7 A lot of further studies investigated the association of other genetic variants of
8 homocysteine metabolism with neuropsychiatric diseases. Although also negative
9 results have been reported, there, e.g., may be associations between the variants
10 CBS c.844_845ins68bp (p.-), MTHFR c.1298A>C (E429A), MS c.2756A>G
11 (D919G), Tc2 c.776C>G (p.R259R), with plasma homocysteine levels, birth defects,
12 cerebrovascular disease, neurodegeneration, (neuro)oncological and psychiatric
13 disorders [54-64]. In the opinion of the authors, none of the genetic variants of
14 homocysteine metabolism has yet been proven to be of sufficient relevance for an
15 individual to justify analysis in the clinical routine beyond studies. Data on the clinical
16 relevance in terms of consequences for therapy or (secondary) prevention of
17 neuropsychiatric diseases may be achieved in the next years.

18 19 **2. Vitamin deficiency in children, adults, and the elderly**

20 21 **2.1. Children**

22 There is a need for a considerable amount of one-carbon groups for brain
23 proliferation, cerebral maturation and myelination, particularly in newborns, children
24 and adolescents during their growth period. Thus, disturbances of folate, vitamin B12
25 and homocysteine metabolism can lead to psychomotor retardation and to a variety

of unspecific neuropsychiatric symptoms [31,32,65-67]. Dysfunctional homocysteine metabolism caused by genetic deficiencies leads to greatly elevated homocysteine concentrations in the plasma ($>100 \mu\text{mol/L}$), often presenting with neuropsychiatric disorders [67-69]. Examples are CBS deficiency [68,69], disturbances of intracellular cobalamin metabolism (cblC, cblD, cblF, cblE-, cblG-defect) and MTHFR deficiency [67]. These rare congenital metabolism disorders need to be dealt with by a specialist department and are not the subject of the present review.

However, also nutritional conditions can lead to severe disturbances of folate, vitamin B12 and homocysteine metabolism in children. In particular, babies breastfed by mothers with vitamin B12 deficiency, e.g., due to vegan diet, can be affected by serious and irreversible CNS damage [70,71].

2.2. Adults

In adults, folate or vitamin B12 deficiency mostly develop for months and years, before e.g. disturbed DNA synthesis and disturbed methylation lead to symptoms of (megaloblastic) anemia and neurological impairment such as forgetfulness, sleeplessness, tiredness, irritability, lethargy and mood swings [72]. Further progression may involve cerebral demyelination, seizures, impairment of the peripheral nervous system like hypo- or paraesthesiae, pareses, depression and dementia [73-75]. Importantly, up to 30% of patients with vitamin B12 deficiency and normal folate levels exclusively show neurological symptoms [75]. In particular, diagnostic attention must be paid to risk groups of folate and vitamin B12 deficiency such as pregnant women, patients with inflammatory gastric conditions, people taking relevant drugs and alcoholics [76].

Vitamin B6 acts as a cofactor in more than 100 enzymatic reactions and is involved in the synthesis of various neurotransmitters [77] such as those occurring in the tryptophan-serotonin metabolism. Vitamin B6 deficiency was suggested to be associated with migraine, chronic pain, seizures and depression [78] as well as cardio-vascular diseases [79], but according to our knowledge there is currently no robust data supporting the idea that vitamin B6 deficiency is epidemiologically relevant for neuropsychiatric diseases in Western European Countries.

2.3 Elderly population

The prevalence of vitamin B12 and folate deficiency increases with age and is common in the elderly [47,80]. Whilst serum and CSF concentrations of folate and vitamin B12 fall, those of homocysteine rise with age [81]. The cause of vitamin deficiency in higher age has been variously ascribed to chronic illnesses, side effects of medications, malabsorption, increased demand and poor diet [82]. In elderly Europeans, the average intake of folate is clearly below the recommended daily dose of 400 µg/day [83].

Many neuropsychiatric diseases connected with the homocysteine metabolism, such as cognitive impairment, dementia, Parkinson's disease and polyneuropathy, have their highest prevalence in elderly persons. Therefore, disturbances of homocysteine metabolism that are associated with neuropsychiatric diseases may be of pronounced importance for the elderly.

3. Neuropsychiatric diseases

The pathophysiological mechanisms of neuropsychiatric diseases can be divided into disease-specific and disease-non-specific damage. Interactions between

mechanisms of neuropsychiatric diseases with folate, vitamin B12 and homocysteine, which have been suggested in the literature, are summarised in table 1.

3.1. Cerebral ischaemia

There is evidence that elevated plasma levels of homocysteine can affect the endothelium, promote growth of (vessel) smooth muscle cells and activate haemostasis [84] (see figure 2). Elevated plasma homocysteine is a confirmed risk factor for atherosclerotic diseases and thromboembolic events [76]. Meta-analyses have proven the association of elevated homocysteine levels with the risk of cerebral ischaemia [53,85,86]. The data collected in 30 retro- and prospective studies showed that a homocysteine difference of $-3 \mu\text{mol/L}$ ($\sim 25\%$) is associated with an approximate 19-24% lower risk of cerebral ischaemia [53,87]. An exponential increase of risk for cerebral ischaemia was shown prospectively for lower dietary folate intake as well as for increased homocysteine levels [88,89]. Each $\mu\text{mol/L}$ of homocysteine led to a risk increase of around 6-7% [90] with a $5 \mu\text{mol/L}$ rise in homocysteine levels raising the risk by 65% [87].

Further, the resulting potential to lower the risk of developing ischaemia by approximately 19% [53,87] was confirmed in randomised controlled therapy studies involving high-risk populations. Although results of the first intervention studies have been regarded as disappointing at first, the Vitamin Intervention for Stroke Prevention (VISP) study based on 3680 probands in whom cerebral ischaemia had occurred, the relative risk of having another ischaemia event was lowered by 21% after 2 years of therapy with 2.5mg folic acid and other B vitamins daily [91].

Substitution of 2.5mg folate/ day together with vitamin B6 and B12 lowered the

relative risk of stroke by 24% during 5 years in the Heart Outcomes Prevention Evaluation (HOPE)-2 study involving 5222 patients with vascular disease or diabetes [92,93]. However, within the NORVIT study, a daily dose of a lower dose of folic acid, i.e., 0.8mg per day, did not prove to be effective in preventing stroke in patients with myocardial infarction [94].

According to a meta-analysis including 16,841 participants from eight randomised studies, folic acid therapy lowered the relative risk of suffering cerebral ischaemia by 18% in all participants [86]. The effect correlated with the observed decrease of homocysteine levels. Importantly, analysis of subgroups showed that the effect of folate on the risk of cerebral ischemia was significant only for participants treated and followed-up for at least 36 months, whereas there was no significant effect in shorter treatment regimens indicating that intervention should be permanent and intervention studies should exceed duration of 36 months.

Food fortification with folic acid in the US and Canada was shown to be associated with a significant drop in the number of deaths by cerebral ischaemia [95].

In summary, folate supplementation has been proven to be effective in primary and secondary prevention of cerebral ischemia. Whether the effect of folate on homocysteine levels or other folate-dependent effects are the underlying mechanisms, remains unknown. The results of further, currently ongoing intervention studies may provide additional information in the near future [96,97].

→ In conclusion, there is evidence that long-term supplementation of folate is effective in secondary stroke prevention (all patients) as well as for primary stroke prevention in patients at risk such as patients with diabetes mellitus or after

1 *myocardial infarction. The best dose or combination with other B-vitamins, if any, has*
2 *not been evaluated yet, but a daily dose of 2.5mg folic acid was shown to be*
3 *effective in the HOPE2 and the VISP study [evidence category IA].*
4
5

6 **3.2. Impairment of cognitive functions**

7

8 Alzheimer's Dementia (AD) is the most frequent form of dementia followed by
9 vascular and mixed (Alzheimer and vascular) dementia [98]. If AD onset could be
10 delayed five years, the number of people affected would approximately halve [99].
11 Folate, vitamin B12 and homocysteine metabolism may have impact on both AD and
12 vascular dementia. Similar to the association of elevated homocysteine levels with
13 major stroke, elevated homocysteine levels are associated with cerebral
14 microangiopathy and microvascular brain lesions as biological correlate of vascular
15 dementia [100-102]. However, there are putative biological mechanisms for the
16 association of homocysteine with dementia exceeding cerebrovascular disease. First
17 of all, homocysteine has neurotoxic effects in cell culture and in animal experiments
18 [9,103]. Possible mechanisms of neurotoxicity of homocysteine are multiple and
19 include activation of NMDA receptors, DNA damage [104], and binding of copper
20 and concomitant cytochrome C oxidase deficiency [105]. Therefore, the presence of
21 elevated homocysteine brain levels may well promote neuronal cell death during
22 neurodegenerative processes.
23 The AD brain is characterised by extracellular beta-amyloid (A β) deposition and
24 intracellular neurofibrillary tangles. Homocysteine metabolism influences the
25 progression of these two histological hallmarks of Alzheimer disease in experimental

models. The amount of A β production depends on expression of the amyloid precursor protein (APP) and different secretases splitting APP in an amyloidogenic or non-amyloidogenic manner [106]. One of the most important mechanisms causing alteration of gene expression is disturbed DNA methylation. DNA methylation in general is accomplished through the specific enzymes, the DNA methyltransferases, which transfer a methyl group to the cytosine of CpG dinucleotides, and the degree of promoter gene CpG methylation is an important factor in gene silencing [107]. Due to the role of SAM as the ubiquitous methyl group donor and SAH as a strong inhibitor of SAM-dependent transmethylation reactions, lower levels of SAM and higher levels of SAH result in a reduced methylation capacity in general and in reduced DNA methylation in particular [108,109]. It has previously been shown that low SAM levels are associated with decreased DNA-demethylation followed by increased expression of presenilin 1 and β -secretase (amyloidogenic pathway), leading to an increase in A β production. Supplementation of SAM prevented these changes in cell culture experiments and mouse models [110-112].

In AD, neurofibrillary tangles are thought to result from hyperphosphorylation of tau protein. High concentrations of hyperphosphorylated tau protein (P-tau) predict the development of dementia [113]. Tau is dephosphorylated by protein phosphatase 2A (PP2A), and methylation of PP2A is required for correct binding and dephosphorylation of tau [114,115]. Incubation of Neuro-2a cells with SAH is associated with a decrease in PP2A methylation and associated with enhanced tau phosphorylation, and PP2s methylation becomes down-regulated in the brains of hyperhomocysteinemic mice [116].

1 Thus, low plasma, CSF and brain levels of SAM and high levels of the
2 methyltransferase-inhibitor SAH may promote both A β production and P-tau
3 accumulation. In support of this hypothesis, SAM levels are decreased in brain tissue
4 and cerebrospinal fluid of Alzheimer patients, [117,118] and increased SAH levels in
5 brain tissues of Alzheimer patients correlate with disease progression and cognitive
6 impairment [119]. Recent findings showed that oral substitution of SAM results in an
7 increase in its plasma and CSF levels, and SAM substitution may lead to some
8 clinical improvement in AD patients according to unconfirmed results [120].

9
10 Another link between homocysteine metabolism and neurodegenerative disorders in
11 general and Alzheimer disease in particular is oxidative stress [121]. First of all, the
12 brain is an organ with a limited baseline transsulfuration capacity, due to the limited
13 activity of CBS and γ -cystathionase [122-124], which is crucial for the transsulfuration
14 reaction of homocysteine to the glutathione component cysteine. Furthermore, the
15 transsulfuration reaction is activated by SAM [125,126]. A lack of brain SAM may
16 result in a reduced antioxidative capacity and increased oxidative stress (figure 1).

17
18 Patients with dementia and reduced memory show lower levels of folate and vitamin
19 B12 and higher levels of homocysteine in plasma and cerebrospinal fluid [127-134].
20 Homocysteine levels are associated with the severity of cognitive, physical and
21 social impairments in demented patients [134-138]. Low folate or high homocysteine
22 levels were reported to be associated with atrophy of the brain, in particular with the
23 cortex, the amygdale and the hippocampi [139-144]. Otherwise apparently healthy
24 people with folate or vitamin B12 deficiency are at elevated risk to develop cognitive
25 impairment and dementia [131,143,145-151]. Differences in homocysteine plasma

1 levels were suggested to explain 5 to 16% of the variance of cognitive function of
2 healthy individuals [149,152]. Elevated homocysteine levels were shown
3 prospectively to be an independent risk factor for mild cognitive impairment (MCI)
4 and its conversion into Alzheimer's disease, and this association is dose-dependent
5 [143,147-149,153-155]. The OR for the risk to develop dementia is increased by a
6 factor of 2.8 to 4.6 in people with homocysteine levels of ≥ 14 $\mu\text{mol/L}$ (compared with
7 <10 $\mu\text{mol/L}$), and a 5 μmol rise in homocysteine levels is associated with a risk
8 increase of approximately 40% [143,149,155]. Hence, a rise in homocysteine levels
9 precedes the clinical onset of dementia, and persons with chronically elevated
10 values have the highest risk of developing dementia. The available data cannot
11 exclude that the rise of homocysteine levels in subjects developing dementia occurs
12 after subclinical disease pathology onset. However, the association of low vitamin
13 B12 and folate levels, which are associated with elevated homocysteine plasma
14 levels, suggests that low folate, low vitamin B12 or homocysteine plasma levels are
15 causally related with dementia.

16 In prospective studies using imaging procedures, it has been proven that both in
17 subjects suffering from dementia and in healthy individuals, low folate and increased
18 homocysteine values are prospectively associated with decreased cortical and
19 hippocampal volume and, in subjects with dementia, also with faster disease
20 progression [139,142,143].

21
22 An increased intake of folate or folic acid (diet or supplementation) was associated
23 with a 50% lower risk of developing AD within 6 years [156]. Although literature is not
24 univocal, some papers reported improved cognitive function in association with
25 administration of folic acid or B vitamins. The largest study conducted so far involved

818 not demented, 50-70 years old participants with hyperhomocysteinemia, but normal vitamin B12 levels. The participants in the treatment group (0.8mg of folic acid over 3 years) performed significantly better in tests of sensomotoric speed, information processing speed and complex memory tasks) than those in the placebo group [157].

There are some limitations concerning the transference of these results to the clinical practise. The studies differed concerning population criteria, vitamin dose, treatment duration and specification of cognitive measurements. In regard of several negative results, further and long term studies are necessary to reconfirm previous positive study results and establish optimal protocols of folate and vitamin B12 intake [158].

The possible protective effect of folate or vitamin B12 against dementia can be expected to be greater the earlier therapy is started, if homocysteine levels are elevated before treatment, and if the duration of therapy is sufficiently long [151,156,157,159,160].

-> We conclude that there is evidence that low levels of folate and vitamin B12 and high levels of homocysteine are risk factors for mild cognitive impairment and dementia. In addition, they dispose towards an unfavourable clinical course. Supplementation of folate and vitamin B12 has beneficial effects for patients with mild cognitive impairment or Alzheimer's disease with evidence of category IIa. One may speculate that B vitamin supplementation might also be beneficial in the maintenance of cognitive function of healthy elderly individuals with high homocysteine levels.

3.3 Depression

Depression is the most frequent psychiatric disease. It is underdiagnosed and undertreated, particularly in older patients [161]. Approximately one third of depressive patients show low levels of folate and elevated levels of homocysteine in serum or erythrocytes, and folate and SAM have been observed to be decreased in the cerebrospinal fluid [162,163]. Accordingly, neuropsychiatric disorders are frequent in patients with megaloblastic anaemia due to folate deficiency [164].

In subjects with severe folate deficiency in the cerebrospinal fluid, greatly altered levels of monoamine metabolites such as hydroxyindole acetic acid and homovanillic acid have been found [165]. Folate also affects the synthesis rate of tetrahydrobiopterin (BH₄), a cofactor for the hydroxylation of phenylalanine and tryptophan and is, therefore, directly involved in the synthesis of monoamine neurotransmitters [166]. Vitamin B6 (pyridoxal-5'-phosphate; PLP) acts as cofactor in the metabolism of tryptophan and serotonin, and a deficiency could be associated with depressive symptoms [78]. However, alternatively, depression may lead to altered nutritional behaviour as the reason for changed folate levels in depressive patients. However, the efficacy of a drug-based treatment using antidepressants seems to be influenced by initial folate values, as treatment is less effective in presence of folate deficiency (delayed or weaker effect) [167-170]. Pre-treatment with folic acid [168] as well as the simultaneous administration of folic acid and fluoxetine leads to a significantly improved efficacy that correlates with changes in homocysteine levels [171], and low folate levels increase the risk of suffering a depressive relapse during fluoxetine therapy [172]. These associations were

1 stronger in women [173,174]. In a large sample of more than 5000 women 20-34
2 years old Kendrick et al found folate levels to be associated with anxiety and
3 depression, but adjustment for socioeconomic and lifestyle factors weakened the
4 association considerably. The authors concluded that socioeconomic factors are
5 much more important for depression in their population of young female participants
6 [175]. However, even in young individuals, this does not rule out a relevant role of
7 folate in case of low folate levels [175] or when moderately decreased folate levels
8 are combined with additional risk factors like the TT genotype of MTHFR c.677C>T
9 [176]. In addition to fluoxetine, folate levels also seem to influence the response to
10 treatment with other antidepressants such as imipramine [167,168], nortriptyline and
11 sertraline [168,177]. Folic acid and SAM may improve the efficacy of an
12 antidepressant-based therapy through better availability of neurotransmitters and via
13 further methylation reactions in the nervous system [178]. Folate and vitamin B12
14 are necessary for SAM synthesis (figure 1). This relation may be a biological basis of
15 the associations observed between folate and depression, as SAM is necessary for
16 the synthesis of dopamine, noradrenaline, serotonin and 5-hydroxyindole-3-acetic
17 acid in the brain [179,180]. Oral and intravenous SAM administration showed a
18 mood-enhancing effect [181] and its antidepressant effect may be comparable with
19 the classical tricyclic antidepressants [182]. However, currently, the clinical use of
20 SAM itself for depression is not approved in most countries.

21
22 Although several categories of antidepressants are available, approximately 30–40%
23 of patients suffer from depression refractory to therapy [183]. Therapy with folic acid
24 should be taken into account as adjunctive treatment and may be beneficial even in
25 cases with normal blood folate levels [184,185].

→ Concerning the majority of published results, folate deficiency favours depression and affects its duration and degree of clinical severity. Folic acid has antidepressant characteristics of its own and improves the therapeutic efficacy of antidepressant drugs. In cases of actual folate deficiency, folic acid administration can be particularly effective [evidence category IA]. In addition, SAM may have antidepressant effects (evidence category IIA).

3.4. Parkinson's Disease and L-DOPA Therapy

Prospective studies in healthy subjects showed no connection between intake of folate and vitamin B12 and the risk of developing Parkinson's disease [186], although one study reported an association of low levels of vitamin B6 and Parkinson's disease [187]. The elevated homocysteine values observed in patients with Parkinson's disease are due to L-DOPA therapy [188]. A large part of the administered L-DOPA receives methyl groups from SAM through the action of catechol-O-methyltransferase (COMT) being converted into 3-O-methyldopa. This reaction leads to a drop in SAM as well as an increase in SAH and homocysteine levels (the latter by 60–80%) [189-192]. Hence, administration of L-DOPA results in a rise in homocysteine values [193,194]. The group with the highest homocysteine concentrations also showed the highest concentrations of 3-OMT [194]. These changes in Parkinson patients can plausibly be reduced with COMT inhibitors [195]. Also in Parkinson patients, elevated homocysteine values in the plasma are

1 associated with a higher incidence of depression [189,196] and an increased risk of
2 cardiovascular disease [188] and cerebral ischaemia [197,198]. The risk of
3 Parkinson patients to develop dementia is four to six times higher [199] and is
4 increased by the presence of hyperhomocysteinemia [196,200,201].

5
6 → *There is no evidence that disturbances of folate, vitamin B12 or homocysteine*
7 *metabolism are relevant for the etiology of Parkinson's disease. However, elevated*
8 *homocysteine levels secondary to L-DOPA treatment in patients with Parkinson's*
9 *disease may accelerate neurodegeneration and may dispose to vascular disease*
10 *and, thus, should be treated with folate and vitamin B12. In addition, vitamin B6 may*
11 *be included in the supplementation regimen as it is necessary for homocysteine*
12 *transsulfuration supporting the synthesis of glutathione. Glutathione is used for the*
13 *defence against oxidative stress that is supposedly involved in neurodegeneration*
14 *such as that seen in Parkinson's disease [evidence category IIb].*

17 **3.5. Schizophrenia**

18
19 Compared with healthy individuals, patients with schizophrenia may have higher
20 plasma homocysteine values [202], and elevated homocysteine or low folate levels
21 seem to correlate with extrapyramidal motor symptoms induced by neuroleptic
22 therapy and with negative symptoms of schizophrenia [203], although these results
23 have not been sufficiently confirmed, yet. A meta-analysis of risk alleles for
24 schizophrenia suggested that the T-allele of MTHFR c.677C>T, the most prevalent
25 genetic disposition for elevated homocysteine plasma levels, may be associated with

schizophrenia [204]. However, the therapy studies published so far did not provide evidence that vitamin supplementation and homocysteine lowering had beneficial effects in the treatment of schizophrenia. A vitamin B6 supplementation was found to lower homocysteine levels, but there was no significant effect on schizophrenia symptoms [205]. A decrease in homocysteine levels in patients with initial values of > 15 µmol/L has allegedly led to a significant improvement of clinical symptoms after three months in a small study [206].

→ *The current data is insufficient to decide whether folate, vitamin B6, vitamin B12 and homocysteine metabolism may be relevant for schizophrenia incidence, clinical course or treatment.* [evidence category III].

3.6 Bipolar disorders

Recent studies observed disturbances of homocysteine metabolism also bipolar disorders (BD). Ozebek et al. showed that homocysteine levels are higher, and folate levels are lower in patients with a bipolar disorder in comparison to healthy controls [207]. Additionally, the variants MTHFR c.1298A>C and MTR c.2756 A>G were shown to be associated with bipolar disorders [207,208]. Hyperhomocysteinemia may also play a role in the pathophysiology of neurocognitive deficits in BD, with a higher impact in older patients, or in patients, who had a delayed onset of illness [209-211]. Prospective studies are required to further analyse the role of homocysteine metabolism in the pathophysiology of bipolar disorders.

→ The current data is insufficient to decide whether folate, vitamin B6, vitamin B12 or homocysteine metabolism may be relevant for bipolar disorder incidence, clinical course or treatment. [evidence category III].

3.7. Multiple Sclerosis

A connection between vitamin B12 deficiency and multiple sclerosis (MS) has been suspected due to the fact that the illness is frequently accompanied by macrocytosis [212]. Irrespective of the course or stage of the disease, elevated concentrations of homocysteine have been found in the plasma and cerebrospinal fluid of MS patients [213,214] as well as lowered concentrations of vitamin B12 [215], whilst folate levels are normal [213]. As alternative to vitamin deficiency or at different stages of disease and treatment, elevated homocysteine levels in multiple sclerosis may also be the consequence of chronic inflammation processes or of cellular immune activation [213,216,217]. Prospective studies on vitamin concentrations in blood and the cerebrospinal fluid before disease onset are lacking. Based on experimental tests, there is a possibility that homocysteine metabolism plays a part in the creation or maintenance of the chronic inflammation process in MS patients. In the animal model of MS, an inhibition of transmethylation suppresses the CD4 cell-mediated autoimmune reactions [218].

Weekly vitamin B12 injections (i.m.) in 138 patients led to an improved clinical picture in the therapy group after 24 weeks whilst a combination therapy mixing vitamin B12 with lofepramine and L-phenylalanine did not produce an additional

therapeutic effect [219]. In a very small therapy group that received multivitamin preparation, the neurological findings were judged to have improved in comparison with the control subjects [220].

→ *There is some evidence that folate, vitamin B12 and homocysteine metabolism is changed in patients with multiple sclerosis. Vitamin B12 and multivitamin substitution have been reported to be beneficial in single studies. However, the current data are insufficient to allow recommendations for diagnosis or treatment to be made [evidence category III].*

3.8. Epilepsy and antiepileptic therapy

Epileptic seizures are caused by the pathological stimulation or a lack of stimulation of nerve cells. NMDA receptors play an important part in the generation and maintenance of epileptic seizures. Homocysteine and other sulphur-containing metabolites (cysteine, homocysteine acid etc.) can trigger epileptic fits as agonists for NMDA receptors [221].

Several anticonvulsive drugs, even those of the newer generation, can interfere with folate metabolism [222]. Their intake lowers folate levels and is therefore associated with an increase in homocysteine concentrations [223]. Changes to plasma homocysteine levels have been observed above all during the application of phenobarbital, carbamazepine, primidone, phenytoin and valproate. Phenytoin and carbamazepine increase homocysteine by lowering folate and vitamin B6 levels [223,224]. Phenytoin, carbamazepine, phenobarbital and primidone induce the cytochrome (CYP) P450 enzyme system in the liver. Phenytoin is a substrate of CYP

1 2C9 as well as 2C19 and inhibits CYP 3A4, 5 and 7. Carbamazepine and
2 phenobarbital induce CYP 2C19 or 2B6, whilst primidone is a substrate of CYP 2C19
3 and does not have an inhibitory effect. In animal trials, phenytoin inhibits MTHFR
4 activity [225]. Disturbed folate metabolism observed during the administration of
5 antiepileptic drugs particularly during the first trimester could be one mechanism for
6 the teratogenicity of some anticonvulsive drugs [226]. In particular, carbamazepine
7 and valproic acid were found to be associated with lowered serum folate levels as a
8 possible risk of neural tube defects, but sufficient data on several antiepileptic drugs
9 are missing [227,228]. This is why folic acid supplementation for women of
10 childbearing age, who are on antiepileptics, is particularly indicated and is urgently
11 recommended by the specialist societies [229,230].

12 Further, elevated homocysteine and lowered folate concentrations can represent a
13 risk factor for the occurrence of an interictal (occurring between seizures) psychosis
14 [231]. Development of depressive symptoms has also been reported by patients on
15 antiepileptic drugs [169,232]. In the animal model, adding folic acid supplements to
16 anticonvulsive therapy has a mood-enhancing effect, improves cognitive functions
17 and raises the seizure threshold [233].

18

19 In children [234] and adults [235] on anticonvulsive drugs, the daily intake of 0.4 to
20 1.0 mg of folic acid lowers elevated homocysteine values in most cases within 1-3
21 months. The treatment of folate-deficient epileptics with 5 mg folic acid over 1–3
22 years was reported to lead to improved initiative, attention, concentration, mood and
23 social behaviour [236]. An adjuvant therapy with vitamin B6 lowered the severity of
24 epileptic seizures even in those forms of paediatric epilepsy that do not belong to the
25 congenital pyridoxine- and pyridoxal-5'-phosphate-dependent forms [77].

→ Antiepileptic drugs can raise homocysteine concentrations supposedly due to their interaction with the folate metabolism. Since valid data are missing for several of the antiepileptic drugs, it is recommended that monitoring of folate and vitamin B12 levels should be performed in all patients treated with antiepileptic drugs in the long-term. In case of abnormal results and, concerning folate, in women of childbearing age, vitamins should be substituted [evidence category IIb and Ic-Ia for women of childbearing age].

4. Safety of multivitamin therapy

Multivitamin therapy may be not safe for everyone. Folate therapy can have negative effects in persons with subclinical vitamin B12 deficiency [237]. This should be avoided by excluding vitamin B12 deficiency before folate therapy, or by supplementing folate together with vitamin B12 in a dose of least 400µg per day. Folate is essential for nucleotide synthesis and DNA methylation. Therefore, folate deficiency has been associated with increased risks of several cancer types. However, high amounts of folate intake may also increase liability to cancer consistent with the role of folate in cell proliferation [238]. Although a safe upper limit of folic acid intake of 1 mg/d for adults and 300–800 g/d for children, depending on age, has been proposed there is no consensus about what blood concentrations of folate might cause harm, if any [239].

Expert commentary

Elevated plasma homocysteine levels are a risk factor for stroke. Whereas, in the NORVIT study, daily substitution of 0.8mg folate did not prove to be effective in

1 secondary stroke prevention, 2.5mg folate significantly reduced the risk for
2 secondary stroke in the HOPE2 and the VISIP study. Accordingly, the meta-analysis
3 of Wang and co-workers which included 8 intervention studies reported a significant
4 preventive effect of folate supplementation against stroke. Most current guidelines,
5 however, do not recommend vitamin substitution for homocysteine lowering in
6 secondary stroke prevention, which supposedly means that some avoidable strokes
7 are not prevented.

8 Elevated levels of plasma and CSF homocysteine may promote diseases like mild
9 cognitive impairment, Alzheimer's disease and vascular dementia by several
10 mechanisms. In addition, homocysteine as NMDA-receptor agonist may directly
11 interfere with antidementive drugs like NMDA-receptor antagonists. In patients with
12 cognitive deficits, folate, vitamin B12 and homocysteine should be determined, and
13 abnormal levels should be treated. Substitution may be beneficial even in cases of
14 values within the reference ranges.

15 Although a lack of folate and vitamin B12 and elevated levels of homocysteine
16 promote neurodegeneration, there is no evidence at the clinical level that folate,
17 vitamin B12 and homocysteine metabolism is involved in the specific aetiology or
18 pathogenesis of Parkinson's disease. However, L-DOPA therapy leads to elevated
19 homocysteine levels which may speculatively have adverse effects on Parkinson's
20 disease in addition to the general risks associated with elevated homocysteine
21 levels. Thus, measurement of plasma levels of homocysteine is recommended in L-
22 DOPA treated patients, and elevated levels should be treated. Similarly, several
23 antiepileptic drugs are associated with decreased levels of folate and increased
24 levels of homocysteine, and they should also be treated. In addition, women of
25 childbearing age treated with antiepileptic drugs should control folate and vitamin

B12 plasma levels and should, if levels are below the reference range, increase the folate prophylaxis for pregnancy to 2.5mg per day starting three months before possible conception in addition to supplementation of vitamin B12, e.g. 100µg per os per day, if necessary [240,241]. Women with instable epilepsy treated with phenytoin should consult a specialist prior to vitamin B12 substitution, as vitamin B12 supplementation may reduce phenytoin efficacy according to this paper's authors' experience.

Depression is associated with low plasma levels of folate and high plasma levels of homocysteine. A reduced synthesis of SAM, which is related to such laboratory findings, may be an underlying mechanism of such associations, as SAM is necessary for neurotransmitter synthesis and may be relevant for the efficacy of several antidepressant drugs. We conclude that there is a case for determination of folate, vitamin B12 and homocysteine levels in patients suffering from depression, and abnormal levels should be treated. Even in patients with normal values, folate substitution may have antidepressant effects or may increase the effects of antidepressant drugs, especially in patients with otherwise drug-resistant depression. Thus, adjuvant treatment with folate is a considerable safe therapeutic option for several patients with depression.

Concerning other neuropsychiatric diseases such as multiple sclerosis or schizophrenia, there is speculation, but no evidence, that folate, vitamin B12 and homocysteine metabolism may have any impact.

Safety: Even after the prolonged use of high doses, the toxicity of folic acid remains low [242]. However, due to the risk of masking megaloblastic anaemia and irreversible neurological disorders, it is not recommended to carry out folic acid therapy without first excluding a causal vitamin B12 deficiency or co-supplementing

1 vitamin B12, in particular in elderly people [242]. In addition, folate supplementation
2 is controversially discussed to promote tumour development or growth. Thus,
3 therapeutic folate supplementation should be restricted to selected populations, until
4 such issues have been solved. It is the opinion of the authors of this paper that folate
5 supplementation is advisable for prevention of stroke in populations at risk, in
6 patients with mild cognitive impairment or dementia and in selected patients suffering
7 from depression in addition to persons with folate levels below the reference range
8 and women who might become pregnant. Based on much therapeutic experience,
9 vitamin B12 (cyanocobalamin and hydroxocobalamin) is considered to be well
10 tolerated. Accordingly, the Food and Nutrition Board of the Institute of Medicine at
11 the National Academy of Sciences has not issued an upper limit for vitamin B12
12 intake. However, high doses exceeding 900µg per os per day may lead to vitamin-
13 B12-acne in rare cases in the experience of the authors.

14 Vitamin B6 is regarded as safe within the dosage range of 2-25 mg in the experience
15 of the authors, which should be sufficient for treatment of levels below the reference
16 ranges [76]. At doses over-exceeding 25 mg, or the more, exceeding 50 mg, side
17 effects like paraesthesia were frequently reported by the respective patients.

18 Concerning laboratory analysis of the vitamins involved in homocysteine metabolism,
19 the determination of the biologically available form of vitamin B12,
20 holotranscobalamin, may be superior to the determination of total vitamin B12 [243].
21 However, this does not reflect the authors' experience. In questionable cases,
22 determination of methylmalonic acid may be much more informative.

24 **Five-year view**

1 For decades, B-vitamins played an important role in the treatment of neurological
2 diseases, and hyperhomocysteinemia has, for a long time, been seen as an
3 important risk factor for cerebrovascular disease. Today, B-vitamins and
4 homocysteine are often regarded as past topics that have lost their eligibility in the
5 treatment and prevention of neuropsychiatric disease, which may in part be due to
6 over-exaggerated expectations or lobbyism. There is evidence that folate, vitamin
7 B12 and homocysteine metabolism can interact with the aetiology and pathology of
8 neuropsychiatric diseases. Manipulation of this metabolism can be beneficial e.g. in
9 relation to stroke prevention and improved clinical course of dementia or depression.
10 Since simple therapies such as vitamin substitution may have clear positive effects
11 on diseases as mentioned it is opportune to reconsider and change current
12 guidelines basing on the evidence of large studies and meta-analyses published in
13 high-ranked peer-reviewed. In five years, additional studies will have finished and,
14 hopefully, we will be able to recommend more exact doses and protocols for
15 treatment and prevention with folate and vitamin B12 for clearer defined populations.

Key issues (8-10 bullet points summarized in the review)

- Folate substitution is effective in selected primary and in secondary prevention of stroke. A dose of 2.5mg per day per os has been proven to be effective, although lower doses may be effective, too. The combination with vitamin B12 may have additional beneficial effects and may improve the safety of folate supplementation. Low doses of 15µg vitamin B12 per day per os may be sufficient, but also higher doses like 100µg can be expected to be safe. The preventive effects increase when supplementation is long-term or continuous. Patients of countries with folate fortification or with low baseline homocysteine levels have lower benefit.
- Folate, vitamin B12 and homocysteine levels should be determined in patients with dementia or depression and, repeatedly, e.g. once a year, in patients treated with antiepileptic drugs or with L-DOPA. Abnormal levels should be treated.
- Folate itself exerts antidepressant effects in animal models and promotes the effects of other antidepressant drugs, in particularly in patients with otherwise drug resistant depression. Folate supplementation maybe considered in patients with depression.
- Substitution of folate and vitamin B12 may improve cognitive functions even in the absence of folate or vitamin B12 deficiency.
- The costs of treatment with folate and vitamin B12 should be covered by health insurance schemes for prevention of vascular events in patients at risk as well as for cognitive impairment, dementia and depression. Treatment guidelines should be modified.

- 1 • The impact of folate, vitamin B12 and homocysteine metabolism on other
- 2 common neuropsychiatric diseases including multiple sclerosis and
- 3 schizophrenia has not been proven.

Evidence categories refer to therapeutic usefulness.

Classification:

Class I Conditions with regard to which there is evidence and/or general agreement that the procedure or treatment is beneficial, useful, and effective.

Class II Conditions with regard to which there is conflicting evidence and/or a divergence of opinion about the usefulness/efficacy of a procedure or treatment.

Class IIa Weight of evidence/opinion is in favour of usefulness/efficacy.

Class IIb Usefulness/efficacy is less well established by evidence/opinion.

Class III Conditions for which there is evidence and/or general agreement that the procedure/treatment is not useful/effective and in some cases may be harmful.

Level of evidence A Data derived from multiple randomized clinical trials.

Level of evidence B Data derived from a single randomized trial or non-randomized studies.

Level of evidence C Only consensus opinion of experts, case studies, or standard-of-care.





Figure 1. The Homocysteine Metabolism



1

2 Figure 2. Electron micrographs of cerebral arterioles in a CBS +/+ mouse fed control
 3 diet (left) and a CBS +/- mouse fed high-methionine diet (right). Vascular lumina (L)
 4 are oriented toward the bottom. Components of the vessel wall include endothelium
 5 (asterisk), elastin (E), smooth muscle (SM), collagen (closed arrows), and basement
 6 membrane (open arrows). Bar = 1 μ m. Reproduced from the article "Structure of
 7 Cerebral Arterioles in Cystathionine b-synthase deficient mice", Circ Res. 2002 Nov
 8 15;91(10):931-7. With kind permission of Gary Baumbach.




Figure 3. Selected mechanisms of neurotoxicity of homocysteine.

Homocysteine (HCYS) can enter the cell from the extracellular space, but also activate the NMDA receptor. This leads to intracellular increase of Ca^{2+} and accumulation of reactive oxygen species (ROS). In addition, homocysteine itself can increase intraneuronal concentrations of ROS. ROS, e.g., increase the intracellular activation of matrix metalloproteinases (MMPs), increase $\text{NF}\kappa\text{B}$ and induce endoplasmatic reticulum stress and mitochondrial dysfunction, cell swelling and osmolysis of the cell. Homocysteine has inhibitory function at the GABA receptor

1 reducing NAPH oxidase activity and promoting oxidative stress. Furthermore,
2 homocysteine forms toxic complexes with copper which can, e.g., induce DNA
3 damage and can cause reduced activity of copper-dependent enzymes like the
4 superoxide dismutase or the cytochrome C oxidase as part of the mitochondrial
5 respiratory chain. Intracellular oxidative stress and interaction with expression,
6 phosphorylation and activation of neuronal proteins are discussed as mechanisms of
7 the association of elevated homocysteine levels with an increased formation of
8 phospho-Tau and β -Amyloid in Alzheimer's disease.

	Hcy↑	FA↓	B12↓
B- and γ -secretases (then A β ↑)	↑	↑	↑
Reactive (free) oxygen radicals	↑	↑	
Ca ²⁺ concentration (cytosol)	↑	↑	
Glutamate excitotoxicity	↑	↑	
MPTP-induced neurotoxicity	↑	↑	
Iron-(copper-)induced neurotoxicity	↑	↑	
DEW-protein phosphorylation	↑	↑	
ATP concentration	↓	↓	
NMDA activation	↑		
Amyloid β -induced neurotoxicity, oxygen radicals, apoptosis	↑	↑	
ADMA	↑		
NO bioavailability	↓	↓	
MPTP toxicity	↑	↑	
DNA strand breaks	↑	↑	
DNA repair	↓	↓	
Transmethylation	↓		
S-Adenosylmethionine (SAM)	↓	(↓)	(↓)
S-Adenosylhomocysteine (SAH)	↑	(↑)	(↑)
Methyltransferase activities	↓	(↓)	(↓)
Endoplasmic reticulum stress	↑	↑	
Cell cycle (Cyclin B and E)	↑		
Endothelial damage	↑		
Activity of Na ⁺ /K ⁺ -ATPase	↓		
Hypomethylation	↑	↑	
COMT activity	↓		
LDL oxidation, lipid peroxidation	↑		
Inflammation, protease activation (MMP, TNF α)	↑	↑	↑
Mitochondrial complex inhibition	↑		
Synapse dysfunction, degeneration	↑	(↑)	
Myelinisation	↓	↓	↓
Tetrahydrobiopterine synthesis		↓	
Neurotransmitter synthesis	↓	↓	
Microangiopathy, ischaemia, hypoxia	↑		
Presenilin 1/2 protein expression	↑	↑	↑
NTPDase* activity	↑		
5'-Nucleotidase activity	↑		
Respiratory chain complex I activity	↓		
Protein kinase-C activity	↑		
Methylation (e.g. DNA)	↓	↓	↓
Na ⁺ , K ⁺ -ATPase activity	↓		
Protein phosphatase 2A	↓	↓	

1

Tyrosinhydroxylase activity	↓	↓	
NF-KB activation	↑		
mGluR activity	↑		
Glutathione-oxidase activity	↓		
HERP protein expression	↑		
Leukoaraiosis	↑	↑	
GFAP mRNA expression			↑
EGF expression			↓
PARP activation	↑		
Caspase activation	↑		

2 Table 1. Metabolic conditions observed in association with increased homocysteine
3 (Hcys), or decreased folic acid (FA) and vitamin B12 plasma levels

4

5 **NMDA** = *N*-methyl-*D*-aspartate-receptor; **NTPDase**= nucleoside triphosphate
6 diphosphohydrolase; **COMT** = catechol-*O*-methyltransferase, **mGluR** = group 1
7 metabotropic glutamate receptors; **ADMA** = asymmetric dimethylarginine; **NO** = nitric
8 oxide; **HERP** = homocysteine-inducible endoplasmic reticulum stress protein; **MPTP**
9 = 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; **MMP** = metalloproteinases, **TNF α** =
10 tumor necrosis factor α ; **GFAP** = glial fibrillary acidic protein; **EGF** = epidermal growth
11 factor; **PARP** = poly-ADP-ribose-polymerase. ↑ = reinforcement, stimulation, ↑ =
12 reduction, inhibition.

13

References

1. Mental and neurological disorders. World Health Organization fact sheet no. 265 December 2001. *Indian J Med Sci*, 56(1), 25-29 (2002).
2. Andlin-Sobocki P, Jonsson B, Wittchen HU, Olesen J. Cost of disorders of the brain in Europe. *Eur J Neurol*, 12 Suppl 1, 1-27 (2005).
3. Murray CJLL, A.D. The global burden of disease: a comprehensive assesment of mortality and disability from diseases, injuries and risk factors in 1990 and projected to 2020. . In: *Global burden of Disease and Injury series*. Bank., HSoPHobotWHOatW (Ed. (Harvard University Press, Cambridge MA, USA., 1996)
4. Truelsen T, Piechowski-Jozwiak B, Bonita R, Mathers C, Bogousslavsky J, Boysen G. Stroke incidence and prevalence in Europe: a review of available data. *Eur J Neurol*, 13(6), 581-598 (2006).
5. de Bree A, van Dusseldorp M, Brouwer IA, van het Hof KH, Steegers-Theunissen RP. Folate intake in Europe: recommended, actual and desired intake. *Eur J Clin Nutr*, 51(10), 643-660 (1997).
6. Eichholzer M, Tonz O, Zimmermann R. Folic acid: a public-health challenge. *Lancet*, 367(9519), 1352-1361 (2006).
7. Al-Tahan J, Gonzalez-Gross M, Pietrzik K. B-vitamin status and intake in European adolescents. A review of the literature. *Nutr Hosp*, 21(4), 452-465 (2006).

- 1 8. Okun JG, Horster F, Farkas LM *et al.* Neurodegeneration in methylmalonic
2 aciduria involves inhibition of complex II and the tricarboxylic acid cycle, and
3 synergistically acting excitotoxicity. *J Biol Chem*, 277(17), 14674-14680
4 (2002).
- 5 9. Ho PI, Ashline D, Dhitavat S *et al.* Folate deprivation induces
6 neurodegeneration: roles of oxidative stress and increased homocysteine.
7 *Neurobiol Dis*, 14(1), 32-42 (2003).
- 8 10. Zieminska E, Matyja E, Kozłowska H, Stafiej A, Lazarewicz JW. Excitotoxic
9 neuronal injury in acute homocysteine neurotoxicity: role of calcium and
10 mitochondrial alterations. *Neurochem Int*, 48(6-7), 491-497 (2006).
- 11 11. Obeid R, Herrmann W. Mechanisms of homocysteine neurotoxicity in
12 neurodegenerative diseases with special reference to dementia. *FEBS Lett*,
13 580(13), 2994-3005 (2006).
- 14 12. Bisschops RH, van der Graaf Y, Mali WP, van der Grond J. Elevated levels of
15 plasma homocysteine are associated with neurotoxicity. *Atherosclerosis*,
16 174(1), 87-92 (2004).
- 17 13. Selhub J. Public health significance of elevated homocysteine. *Food Nutr Bull*,
18 29(2 Suppl), S116-125 (2008).
- 19 14. Obeid R, McCaddon A, Herrmann W. The role of hyperhomocysteinemia and
20 B-vitamin deficiency in neurological and psychiatric diseases. *Clin Chem Lab*
21 *Med*, 45(12), 1590-1606 (2007).

- 1 15. Fowler B. Homocysteine: overview of biochemistry, molecular biology, and
2 role in disease processes. *Semin. Vasc. Med.*, 5(2), 77-86 (2005).
- 3 16. Chiang PK, Gordon RK, Tal J *et al.* S-Adenosylmethionine and methylation.
4 *Faseb J*, 10(4), 471-480 (1996).
- 5 17. Bottiglieri T, Hyland K, Reynolds EH. The clinical potential of ademetionine
6 (S-adenosylmethionine) in neurological disorders. *Drugs*, 48(2), 137-152
7 (1994).
- 8 18. Axelrod J. Methylation reactions in the formation and metabolism of
9 catecholamines and other biogenic amines. *Pharmacol Rev*, 18(1), 95-113
10 (1966).
- 11 19. McKeever M, Molloy A, Weir DG *et al.* An abnormal methylation ratio induces
12 hypomethylation in vitro in the brain of pig and man, but not in rat. *Clin Sci*
13 (*Lond*), 88(1), 73-79 (1995).
- 14 20. Weir DG, Molloy AM, Keating JN *et al.* Correlation of the ratio of S-adenosyl-
15 L-methionine to S-adenosyl-L-homocysteine in the brain and cerebrospinal
16 fluid of the pig: implications for the determination of this methylation ratio in
17 human brain. *Clin Sci (Lond)*, 82(1), 93-97 (1992).
- 18 21. Smulders YM, Smith DE, Kok RM *et al.* Cellular folate vitamer distribution
19 during and after correction of vitamin B12 deficiency: a case for the
20 methylfolate trap. *Br J Haematol*, 132(5), 623-629 (2006).

- 1 22. Surtees R, Leonard J, Austin S. Association of demyelination with deficiency
2 of cerebrospinal-fluid S-adenosylmethionine in inborn errors of methyl-transfer
3 pathway. *Lancet*, 338(8782-8783), 1550-1554 (1991).
- 4 23. Sunden SL, Renduchintala MS, Park EI, Miklasz SD, Garrow TA. Betaine-
5 homocysteine methyltransferase expression in porcine and human tissues
6 and chromosomal localization of the human gene. *Arch Biochem Biophys*,
7 345(1), 171-174 (1997).
- 8 24. Finkelstein JD. Pathways and regulation of homocysteine metabolism in
9 mammals. *Semin Thromb Hemost*, 26(3), 219-225 (2000).
- 10 25. Vitvitsky V, Thomas M, Ghorpade A, Gendelman HE, Banerjee R. A functional
11 transsulfuration pathway in the brain links to glutathione homeostasis. *J Biol*
12 *Chem*, 281(47), 35785-35793 (2006).
- 13 26. Spector R, Coakley G, Blakely R. Methionine recycling in brain: a role for
14 folates and vitamin B-12. *J Neurochem*, 34(1), 132-137 (1980).
- 15 27. Cobalamin deficiency and neuropsychiatric disorders. *N Engl J Med*, 319(26),
16 1733-1735 (1988).
- 17 28. Kolker S, Ahlemeyer B, Krieglstein J, Hoffmann GF. Methylmalonic acid
18 induces excitotoxic neuronal damage in vitro. *J Inherit Metab Dis*, 23(4), 355-
19 358 (2000).
- 20 29. Kamen BA, Smith AK. A review of folate receptor alpha cycling and 5-
21 methyltetrahydrofolate accumulation with an emphasis on cell models in vitro.
22 *Adv Drug Deliv Rev*, 56(8), 1085-1097 (2004).

- 1 30. Wollack JB, Makori B, Ahlawat S *et al.* Characterization of folate uptake by
2 choroid plexus epithelial cells in a rat primary culture model. *J Neurochem*,
3 104(6), 1494-1503 (2008).
- 4 31. Ramaekers VT, Blau N. Cerebral folate deficiency. *Dev Med Child Neurol*,
5 46(12), 843-851 (2004).
- 6 32. Reynolds EH, Gallagher BB, Mattson RH, Bowers M, Johnson AL.
7 Relationship between serum and cerebrospinal fluid folate. *Nature*,
8 240(5377), 155-157 (1972).
- 9 33. Spector R, Eells J. Deoxynucleoside and vitamin transport into the central
10 nervous system. *Fed Proc*, 43(2), 196-200 (1984).
- 11 34. Spector R, Johanson CE. Vitamin transport and homeostasis in mammalian
12 brain: focus on Vitamins B and E. *J Neurochem*, 103(2), 425-438 (2007).
- 13 35. Sharma P, Senthilkumar RD, Brahmachari V *et al.* Mining literature for a
14 comprehensive pathway analysis: a case study for retrieval of homocysteine
15 related genes for genetic and epigenetic studies. *Lipids Health Dis*, 5, 1
16 (2006).
- 17 36. Gellekink H, den Heijer M, Heil SG, Blom HJ. Genetic determinants of plasma
18 total homocysteine. *Semin. Vasc. Med.*, 5(2), 98-109 (2005).
- 19 37. Frosst P, Blom HJ, Milos R *et al.* A candidate genetic risk factor for vascular
20 disease: a common mutation in methylenetetrahydrofolate reductase.
21 *Nat. Genet.*, 10(1), 111-113 (1995).

- 1 38. Jacques PF, Bostom AG, Williams RR *et al.* Relation between folate status, a
2 common mutation in methylenetetrahydrofolate reductase, and plasma
3 homocysteine concentrations. *Circulation*, 93(1), 7-9 (1996).
- 4 39. Kauwell GP, Wilsky CE, Cerda JJ *et al.* Methylenetetrahydrofolate reductase
5 mutation (677C-->T) negatively influences plasma homocysteine response to
6 marginal folate intake in elderly women. *Metabolism*, 49(11), 1440-1443
7 (2000).
- 8 40. Malinow MR, Nieto FJ, Kruger WD *et al.* The effects of folic acid
9 supplementation on plasma total homocysteine are modulated by multivitamin
10 use and methylenetetrahydrofolate reductase genotypes. *Arterioscler Thromb*
11 *Vasc Biol*, 17(6), 1157-1162 (1997).
- 12 41. Fohr IP, Prinz-Langenohl R, Bronstrup A *et al.* 5,10-Methylenetetrahydrofolate
13 reductase genotype determines the plasma homocysteine-lowering effect of
14 supplementation with 5-methyltetrahydrofolate or folic acid in healthy young
15 women. *Am J Clin Nutr*, 75(2), 275-282 (2002).
- 16 42. de Bree A, Verschuren WM, Bjorke-Monsen AL *et al.* Effect of the
17 methylenetetrahydrofolate reductase 677C-->T mutation on the relations
18 among folate intake and plasma folate and homocysteine concentrations in a
19 general population sample. *Am J Clin Nutr*, 77(3), 687-693 (2003).
- 20 43. Brattstrom L, Wilcken DE, Ohrvik J, Brudin L. Common
21 methylenetetrahydrofolate reductase gene mutation leads to
22 hyperhomocysteinemia but not to vascular disease: the result of a meta-
23 analysis. *Circulation*, 98(23), 2520-2526 (1998).

- 1 44. Klerk M, Verhoef P, Clarke R, Blom HJ, Kok FJ, Schouten EG. MTHFR 677C-
2 ->T polymorphism and risk of coronary heart disease: a meta-analysis. *Jama*,
3 288(16), 2023-2031 (2002).
- 4 45. Harmon DL, Doyle RM, Meleady R *et al.* Genetic analysis of the thermolabile
5 variant of 5, 10-methylenetetrahydrofolate reductase as a risk factor for
6 ischemic stroke. *Arterioscler. Thromb. Vasc. Biol.*, 19(2), 208-211 (1999).
- 7 46. Dierkes J, Jeckel A, Ambrosch A, Westphal S, Luley C, Boeing H. Factors
8 explaining the difference of total homocysteine between men and women in
9 the European Investigation Into Cancer and Nutrition Potsdam study.
10 *Metabolism*, 50(6), 640-645 (2001).
- 11 47. Selhub J, Jacques PF, Wilson PW, Rush D, Rosenberg IH. Vitamin status and
12 intake as primary determinants of homocysteinemia in an elderly population.
13 *JAMA*, 270(22), 2693-2698 (1993).
- 14 48. Norlund L, Grubb A, Fex G *et al.* The increase of plasma homocysteine
15 concentrations with age is partly due to the deterioration of renal function as
16 determined by plasma cystatin C. *Clin Chem Lab Med*, 36(3), 175-178 (1998).
- 17 49. Jacques PF, Bostom AG, Wilson PW, Rich S, Rosenberg IH, Selhub J.
18 Determinants of plasma total homocysteine concentration in the Framingham
19 Offspring cohort. *Am J Clin Nutr*, 73(3), 613-621 (2001).
- 20 50. Anello G, Gueant-Rodriguez RM, Bosco P *et al.* Homocysteine and
21 methylenetetrahydrofolate reductase polymorphism in Alzheimer's disease.
22 *Neuroreport*, 15(5), 859-861 (2004).

- 1 51. Nishiyama M, Kato Y, Hashimoto M, Yukawa S, Omori K. Apolipoprotein E,
2 methylenetetrahydrofolate reductase (MTHFR) mutation and the risk of senile
3 dementia--an epidemiological study using the polymerase chain reaction
4 (PCR) method. *J.Epidemiol.*, 10(3), 163-172 (2000).
- 5 52. Kelly PJ, Rosand J, Kistler JP *et al.* Homocysteine, MTHFR 677C-->T
6 polymorphism, and risk of ischemic stroke: results of a meta-analysis.
7 *Neurology*, 59(4), 529-536 (2002).
- 8 53. Homocysteine and risk of ischemic heart disease and stroke: a meta-analysis.
9 *Jama*, 288(16), 2015-2022 (2002).
- 10 54. Wullner U, Kolsch H, Linnebank M. Methylenetetrahydrofolate reductase in
11 Parkinson's disease. *Ann.Neurol.*, 58(6), 972-973 (2005).
- 12 55. Linnebank M, Linnebank A, Jeub M *et al.* Lack of genetic dispositions to
13 hyperhomocysteinemia in Alzheimer disease. *Am.J.Med.Genet.A*, 131(1),
14 101-102 (2004).
- 15 56. Skibola CF, Smith MT, Kane E *et al.* Polymorphisms in the
16 methylenetetrahydrofolate reductase gene are associated with susceptibility
17 to acute leukemia in adults. *Proc.Natl.Acad.Sci.U.S.A*, 96(22), 12810-12815
18 (1999).
- 19 57. Matsuo K, Hamajima N, Suzuki R *et al.* Methylenetetrahydrofolate reductase
20 gene (MTHFR) polymorphisms and reduced risk of malignant lymphoma.
21 *Am.J.Hematol.*, 77(4), 351-357 (2004).

- 1 58. Keku T, Millikan R, Worley K *et al.* 5,10-Methylenetetrahydrofolate reductase
2 codon 677 and 1298 polymorphisms and colon cancer in African Americans
3 and whites. *Cancer Epidemiol.Biomarkers Prev.*, 11(12), 1611-1621 (2002).
- 4 59. Wang J, Gajalakshmi V, Jiang J *et al.* Associations between 5,10-
5 methylenetetrahydrofolate reductase codon 677 and 1298 genetic
6 polymorphisms and environmental factors with reference to susceptibility to
7 colorectal cancer: a case-control study in an Indian population. *Int.J.Cancer*,
8 118(4), 991-997 (2006).
- 9 60. van der Put NM, Gabreels F, Stevens EM *et al.* A second common mutation in
10 the methylenetetrahydrofolate reductase gene: an additional risk factor for
11 neural-tube defects? *Am.J.Hum.Genet.*, 62(5), 1044-1051 (1998).
- 12 61. Linnebank M, Schmidt S, Kolsch H *et al.* The methionine synthase
13 polymorphism D919G alters susceptibility to primary central nervous system
14 lymphoma. *Br.J.Cancer*, 90(10), 1969-1971 (2004).
- 15 62. Semmler A, Simon M, Moskau S, Linnebank M. The Methionine Synthase
16 Polymorphism c.2756A>G Alters Susceptibility to Glioblastoma Multiforme.
17 *Cancer Epidemiol Biomarkers Prev*, 15(11), 2314-2316 (2006).
- 18 63. Semmler A, Simon M, Moskau S, Linnebank M. Polymorphisms of methionine
19 metabolism and susceptibility to meningioma formation: laboratory
20 investigation. *J Neurosurg*, 108(5), 999-1004 (2008).
- 21 64. Semmler A, Linnebank M, Krex D *et al.* Polymorphisms of Homocysteine
22 Metabolism Are Associated with Intracranial Aneurysms. *Cerebrovasc Dis*,
23 26(4), 425-429 (2008).

- 1 65. Ramaekers VT, Hausler M, Opladen T, Heimann G, Blau N. Psychomotor
2 retardation, spastic paraplegia, cerebellar ataxia and dyskinesia associated
3 with low 5-methyltetrahydrofolate in cerebrospinal fluid: a novel
4 neurometabolic condition responding to folinic acid substitution.
5 *Neuropediatrics*, 33(6), 301-308 (2002).
- 6 66. Ramaekers VT, Rothenberg SP, Sequeira JM *et al.* Autoantibodies to folate
7 receptors in the cerebral folate deficiency syndrome. *N Engl J Med*, 352(19),
8 1985-1991 (2005).
- 9 67. Fowler B. Genetic defects of folate and cobalamin metabolism. *Eur J Pediatr*,
10 157 Suppl 2, S60-66 (1998).
- 11 68. Yap S, Naughten E. Homocystinuria due to cystathionine beta-synthase
12 deficiency in Ireland: 25 years' experience of a newborn screened and treated
13 population with reference to clinical outcome and biochemical control. *J Inherit*
14 *Metab Dis*, 21(7), 738-747 (1998).
- 15 69. Yap S. Classical homocystinuria: vascular risk and its prevention. *J Inherit*
16 *Metab Dis*, 26(2-3), 259-265 (2003).
- 17 70. Lovblad K, Ramelli G, Remonda L, Nirkko AC, Ozdoba C, Schroth G.
18 Retardation of myelination due to dietary vitamin B12 deficiency: cranial MRI
19 findings. *Pediatr Radiol*, 27(2), 155-158 (1997).
- 20 71. Rosenblatt DS, Whitehead VM. Cobalamin and folate deficiency: acquired
21 and hereditary disorders in children. *Semin Hematol*, 36(1), 19-34 (1999).

- 1 72. Rosenberg IH, Miller JW. Nutritional factors in physical and cognitive functions
2 of elderly people. *Am J Clin Nutr*, 55(6 Suppl), 1237S-1243S (1992).
- 3 73. Muller WH, Froscher W. [Neurologic and psychologic disorders in folic acid
4 deficiency]. *Fortschr Neurol Psychiatr*, 57(9), 395-401 (1989).
- 5 74. Savage DG, Lindenbaum J. Neurological complications of acquired cobalamin
6 deficiency: clinical aspects. *Baillieres Clin Haematol*, 8(3), 657-678 (1995).
- 7 75. Lindenbaum J, Healton EB, Savage DG *et al*. Neuropsychiatric disorders
8 caused by cobalamin deficiency in the absence of anemia or macrocytosis. *N*
9 *Engl J Med*, 318(26), 1720-1728 (1988).
- 10 76. Stanger O, Herrmann W, Pietrzik K *et al*. Clinical use and rational
11 management of homocysteine, folic acid, and B vitamins in cardiovascular
12 and thrombotic diseases. *Z Kardiol*, 93(6), 439-453 (2004).
- 13 77. Clayton PT. B6-responsive disorders: a model of vitamin dependency. *J*
14 *Inherit Metab Dis*, 29(2-3), 317-326 (2006).
- 15 78. Hvas AM, Juul S, Bech P, Nexø E. Vitamin B6 level is associated with
16 symptoms of depression. *Psychother Psychosom*, 73(6), 340-343 (2004).
- 17 79. Robinson K, Arheart K, Refsum H *et al*. Low circulating folate and vitamin B6
18 concentrations: risk factors for stroke, peripheral vascular disease, and
19 coronary artery disease. European COMAC Group. *Circulation*, 97(5), 437-
20 443 (1998).
- 21 80. Clarke R, Grimley Evans J, Schneede J *et al*. Vitamin B12 and folate
22 deficiency in later life. *Age Ageing*, 33(1), 34-41 (2004).

- 1 81. Bottiglieri T, Reynolds EH, Laundry M. Folate in CSF and age. *J Neurol*
2 *Neurosurg Psychiatry*, 69(4), 562 (2000).
- 3 82. Reynolds EH. Folic acid, ageing, depression, and dementia. *Bmj*, 324(7352),
4 1512-1515 (2002).
- 5 83. Fabian E, Elmadfa I. Nutritional situation of the elderly in the European Union:
6 data of the European Nutrition and Health Report (2004). *Ann Nutr Metab*, 52
7 Suppl 1, 57-61 (2008).
- 8 84. Domagala TB, Undas A, Libura M, Szczeklik A. Pathogenesis of vascular
9 disease in hyperhomocysteinaemia. *J Cardiovasc Risk*, 5(4), 239-247 (1998).
- 10 85. Casas JP, Bautista LE, Smeeth L, Sharma P, Hingorani AD. Homocysteine
11 and stroke: evidence on a causal link from mendelian randomisation. *Lancet*,
12 365(9455), 224-232 (2005).
- 13 86. Wang X, Qin X, Demirtas H *et al.* Efficacy of folic acid supplementation in
14 stroke prevention: a meta-analysis. *Lancet*, 369(9576), 1876-1882 (2007).
- 15 87. Wald DS, Law M, Morris JK. Homocysteine and cardiovascular disease:
16 evidence on causality from a meta-analysis. *Bmj*, 325(7374), 1202 (2002).
- 17 88. Bazzano LA, He J, Ogden LG *et al.* Dietary intake of folate and risk of stroke
18 in US men and women: NHANES I Epidemiologic Follow-up Study. National
19 Health and Nutrition Examination Survey. *Stroke*, 33(5), 1183-1188 (2002).
- 20 89. Bostom AG, Rosenberg IH, Silbershatz H *et al.* Nonfasting plasma total
21 homocysteine levels and stroke incidence in elderly persons: the Framingham
22 Study. *Ann Intern Med*, 131(5), 352-355 (1999).

- 1 90. Bots ML, Launer LJ, Lindemans J *et al.* Homocysteine and short-term risk of
2 myocardial infarction and stroke in the elderly: the Rotterdam Study. *Arch*
3 *Intern Med*, 159(1), 38-44 (1999).
- 4 91. Spence JD, Bang H, Chambless LE, Stampfer MJ. Vitamin Intervention For
5 Stroke Prevention trial: an efficacy analysis. *Stroke*, 36(11), 2404-2409
6 (2005).
- 7 92. Lonn E, Yusuf S, Arnold MJ *et al.* Homocysteine lowering with folic acid and B
8 vitamins in vascular disease. *N Engl J Med*, 354(15), 1567-1577 (2006).
- 9 93. Saposnik G, Ray JG, Sheridan P, McQueen M, Lonn E. Homocysteine-
10 Lowering Therapy and Stroke Risk, Severity, and Disability. Additional
11 Findings From the HOPE 2 Trial. *Stroke*, (2009).
- 12 94. Bonna KH, Njolstad I, Ueland PM *et al.* Homocysteine lowering and
13 cardiovascular events after acute myocardial infarction. *N Engl J Med*,
14 354(15), 1578-1588 (2006).
- 15 95. Yang Q, Botto LD, Erickson JD *et al.* Improvement in stroke mortality in
16 Canada and the United States, 1990 to 2002. *Circulation*, 113(10), 1335-1343
17 (2006).
- 18 96. Clarke R, Lewington S, Sherliker P, Armitage J. Effects of B-vitamins on
19 plasma homocysteine concentrations and on risk of cardiovascular disease
20 and dementia. *Curr Opin Clin Nutr Metab Care*, 10(1), 32-39 (2007).

- 1 97. Homocysteine-lowering trials for prevention of cardiovascular events: a review
2 of the design and power of the large randomized trials. *Am Heart J*, 151(2),
3 282-287 (2006).
- 4 98. Hofman A, Ott A, Breteler MM *et al.* Atherosclerosis, apolipoprotein E, and
5 prevalence of dementia and Alzheimer's disease in the Rotterdam Study.
6 *Lancet*, 349(9046), 151-154 (1997).
- 7 99. Katzman R. Luigi Amaducci memorial award winner's paper 2003. A
8 neurologist's view of Alzheimer's disease and dementia. *Int Psychogeriatr*,
9 16(3), 259-273 (2004).
- 10 100. Sachdev PS, Valenzuela MJ, Brodaty H *et al.* Homocysteine as a risk factor
11 for cognitive impairment in stroke patients. *Dement Geriatr Cogn Disord*,
12 15(3), 155-162 (2003).
- 13 101. Longstreth WT, Jr., Katz R, Olson J *et al.* Plasma total homocysteine levels
14 and cranial magnetic resonance imaging findings in elderly persons: the
15 Cardiovascular Health Study. *Arch Neurol*, 61(1), 67-72 (2004).
- 16 102. Fassbender K, Mielke O, Bertsch T, Nafe B, Froschen S, Hennerici M.
17 Homocysteine in cerebral macroangiography and microangiopathy. *Lancet*,
18 353(9164), 1586-1587 (1999).
- 19 103. Kruman, II, Kumaravel TS, Lohani A *et al.* Folic acid deficiency and
20 homocysteine impair DNA repair in hippocampal neurons and sensitize them
21 to amyloid toxicity in experimental models of Alzheimer's disease. *J Neurosci*,
22 22(5), 1752-1762 (2002).

- 1 104. Ho PI, Ortiz D, Rogers E, Shea TB. Multiple aspects of homocysteine
2 neurotoxicity: glutamate excitotoxicity, kinase hyperactivation and DNA
3 damage. *J Neurosci Res*, 70(5), 694-702 (2002).
- 4 105. Linnebank M, Lutz H, Jarre E *et al.* Binding of copper is a mechanism of
5 homocysteine toxicity leading to COX deficiency and apoptosis in primary
6 neurons, PC12 and SHSY-5Y cells. *Neurobiol Dis*, 23(3), 725-730 (2006).
- 7 106. Evin G, Zhu A, Holsinger RM, Masters CL, Li QX. Proteolytic processing of
8 the Alzheimer's disease amyloid precursor protein in brain and platelets. *J*
9 *Neurosci Res*, 74(3), 386-392 (2003).
- 10 107. Gopalakrishnan S, Van Emburgh BO, Robertson KD. DNA methylation in
11 development and human disease. *Mutat Res*, 647(1-2), 30-38 (2008).
- 12 108. Finkelstein JD. The metabolism of homocysteine: pathways and regulation.
13 *Eur.J.Pediatr.*, 157 Suppl 2, S40-S44 (1998).
- 14 109. Yi P, Melnyk S, Pogribna M, Pogribny IP, Hine RJ, James SJ. Increase in
15 plasma homocysteine associated with parallel increases in plasma S-
16 adenosylhomocysteine and lymphocyte DNA hypomethylation. *J Biol Chem*,
17 275(38), 29318-29323 (2000).
- 18 110. Scarpa S, Fuso A, D'Anselmi F, Cavallaro RA. Presenilin 1 gene silencing by
19 S-adenosylmethionine: a treatment for Alzheimer disease? *FEBS Lett*, 541(1-
20 3), 145-148 (2003).
- 21 111. Fuso A, Seminara L, Cavallaro RA, D'Anselmi F, Scarpa S. S-
22 adenosylmethionine/homocysteine cycle alterations modify DNA methylation

- 1 status with consequent deregulation of PS1 and BACE and beta-amyloid
2 production. *Mol Cell Neurosci*, 28(1), 195-204 (2005).
- 3 112. Chan A, Shea TB. Folate deprivation increases presenilin expression,
4 gamma-secretase activity, and Abeta levels in murine brain: potentiation by
5 ApoE deficiency and alleviation by dietary S-adenosyl methionine. *J*
6 *Neurochem*, 102(3), 753-760 (2007).
- 7 113. Hansson O, Zetterberg H, Buchhave P, Londos E, Blennow K, Minthon L.
8 Association between CSF biomarkers and incipient Alzheimer's disease in
9 patients with mild cognitive impairment: a follow-up study. *Lancet Neurol*, 5(3),
10 228-234 (2006).
- 11 114. Vafai SB, Stock JB. Protein phosphatase 2A methylation: a link between
12 elevated plasma homocysteine and Alzheimer's Disease. *FEBS Lett*, 518(1-
13 3), 1-4 (2002).
- 14 115. Sontag E, Nunbhakdi-Craig V, Lee G, Bloom GS, Mumby MC. Regulation of
15 the phosphorylation state and microtubule-binding activity of Tau by protein
16 phosphatase 2A. *Neuron*, 17(6), 1201-1207 (1996).
- 17 116. Sontag E, Nunbhakdi-Craig V, Sontag JM *et al*. Protein phosphatase 2A
18 methyltransferase links homocysteine metabolism with tau and amyloid
19 precursor protein regulation. *J Neurosci*, 27(11), 2751-2759 (2007).
- 20 117. Morrison LD, Smith DD, Kish SJ. Brain S-adenosylmethionine levels are
21 severely decreased in Alzheimer's disease. *J. Neurochem.*, 67(3), 1328-1331
22 (1996).

- 1 118. Bottiglieri T, Reynolds EH, Toone BK, Carney MW. CSF S-
2 adenosylmethionine in neuropsychiatric disorders. *Lancet*, 338(8759), 121
3 (1991).
- 4 119. Kennedy BP, Bottiglieri T, Arning E, Ziegler MG, Hansen LA, Masliah E.
5 Elevated S-adenosylhomocysteine in Alzheimer brain: influence on
6 methyltransferases and cognitive function. *J Neural Transm*, 111(4), 547-567
7 (2004).
- 8 120. Shea TB, Chan A. S-adenosyl methionine: a natural therapeutic agent
9 effective against multiple hallmarks and risk factors associated with
10 Alzheimer's disease. *J Alzheimers Dis*, 13(1), 67-70 (2008).
- 11 121. Topal G, Brunet A, Millanvoye E *et al.* Homocysteine induces oxidative stress
12 by uncoupling of NO synthase activity through reduction of
13 tetrahydrobiopterin. *Free Radic Biol Med*, 36(12), 1532-1541 (2004).
- 14 122. Finkelstein JD. Methionine metabolism in mammals. *J Nutr Biochem*, 1(5),
15 228-237 (1990).
- 16 123. Ishii I, Akahoshi N, Yu XN *et al.* Murine cystathionine gamma-lyase: complete
17 cDNA and genomic sequences, promoter activity, tissue distribution and
18 developmental expression. *Biochem J*, 381(Pt 1), 113-123 (2004).
- 19 124. Vitvitsky V, Dayal S, Stabler S *et al.* Perturbations in homocysteine-linked
20 redox homeostasis in a murine model for hyperhomocysteinemia. *Am J*
21 *Physiol Regul Integr Comp Physiol*, 287(1), R39-46 (2004).

- 1 125. Selhub J, Miller JW. The pathogenesis of homocysteinemia: interruption of the
2 coordinate regulation by S-adenosylmethionine of the remethylation and
3 transsulfuration of homocysteine. *Am J Clin Nutr*, 55(1), 131-138 (1992).
- 4 126. Mudd SH, Levy HL, Kraus JP. Disorders of transsulfuration. (Ed.^(Eds)
5 (Scriver CR, Beaudet AL, Sly WS, Valle D, Childs B, Kinzler K, Vogelstein B.
6 Editors, New York: Mc Graw-Hill, 2001) 2007-2056.
- 7 127. Serot JM, Christmann D, Dubost T, Bene MC, Faure GC. CSF-folate levels
8 are decreased in late-onset AD patients. *J Neural Transm*, 108(1), 93-99
9 (2001).
- 10 128. Kristensen MO, Gulmann NC, Christensen JE, Ostergaard K, Rasmussen K.
11 Serum cobalamin and methylmalonic acid in Alzheimer dementia. *Acta Neurol*
12 *Scand*, 87(6), 475-481 (1993).
- 13 129. Lewis MS, Miller LS, Johnson MA, Dolce EB, Allen RH, Stabler SP. Elevated
14 methylmalonic acid is related to cognitive impairment in older adults enrolled
15 in an elderly nutrition program. *J Nutr Elder*, 24(3), 47-65 (2005).
- 16 130. Refsum H, Smith AD. Low vitamin B-12 status in confirmed Alzheimer's
17 disease as revealed by serum holotranscobalamin. *J Neurol Neurosurg*
18 *Psychiatry*, 74(7), 959-961 (2003).
- 19 131. Wang HX, Wahlin A, Basun H, Fastbom J, Winblad B, Fratiglioni L. Vitamin
20 B(12) and folate in relation to the development of Alzheimer's disease.
21 *Neurology*, 56(9), 1188-1194 (2001).

- 1 132. McCaddon A, Regland B, Hudson P, Davies G. Functional vitamin B(12)
2 deficiency and Alzheimer disease. *Neurology*, 58(9), 1395-1399 (2002).
- 3 133. Leblhuber F, Walli J, Artner-Dworzak E *et al.* Hyperhomocysteinemia in
4 dementia. *J Neural Transm*, 107(12), 1469-1474 (2000).
- 5 134. Morris MS. Homocysteine and Alzheimer's disease. *Lancet Neurol*, 2(7), 425-
6 428 (2003).
- 7 135. Lehmann M, Gottfries CG, Regland B. Identification of cognitive impairment in
8 the elderly: homocysteine is an early marker. *Dement Geriatr Cogn Disord*,
9 10(1), 12-20 (1999).
- 10 136. Nilsson K, Gustafson L, Hultberg B. Plasma homocysteine concentration
11 relates to the severity but not to the duration of Alzheimer's disease. *Int J*
12 *Geriatr Psychiatry*, 19(7), 666-672 (2004).
- 13 137. Nilsson K, Gustafson L, Hultberg B. The plasma homocysteine concentration
14 is better than that of serum methylmalonic acid as a marker for
15 sociopsychological performance in a psychogeriatric population. *Clin Chem*,
16 46(5), 691-696 (2000).
- 17 138. Wright CB, Lee HS, Paik MC, Stabler SP, Allen RH, Sacco RL. Total
18 homocysteine and cognition in a tri-ethnic cohort: the Northern Manhattan
19 Study. *Neurology*, 63(2), 254-260 (2004).
- 20 139. den Heijer T, Vermeer SE, Clarke R *et al.* Homocysteine and brain atrophy on
21 MRI of non-demented elderly. *Brain*, 126(Pt 1), 170-175 (2003).

- 1 140. Scott TM, Tucker KL, Bhadelia A *et al.* Homocysteine and B vitamins relate to
2 brain volume and white-matter changes in geriatric patients with psychiatric
3 disorders. *Am J Geriatr Psychiatry*, 12(6), 631-638 (2004).
- 4 141. Sachdev P. Homocysteine, cerebrovascular disease and brain atrophy.
5 *J.Neurol.Sci.*, 226(1-2), 25-29 (2004).
- 6 142. Snowdon DA, Tully CL, Smith CD, Riley KP, Markesbery WR. Serum folate
7 and the severity of atrophy of the neocortex in Alzheimer disease: findings
8 from the Nun study. *Am.J.Clin.Nutr.*, 71(4), 993-998 (2000).
- 9 143. Clarke R, Smith AD, Jobst KA, Refsum H, Sutton L, Ueland PM. Folate,
10 vitamin B12, and serum total homocysteine levels in confirmed Alzheimer
11 disease. *Arch Neurol*, 55(11), 1449-1455 (1998).
- 12 144. Williams JH, Pereira EA, Budge MM, Bradley KM. Minimal hippocampal width
13 relates to plasma homocysteine in community-dwelling older people. *Age*
14 *Ageing*, 31(6), 440-444 (2002).
- 15 145. Kado DM, Karlamangla AS, Huang MH *et al.* Homocysteine versus the
16 vitamins folate, B6, and B12 as predictors of cognitive function and decline in
17 older high-functioning adults: MacArthur Studies of Successful Aging. *Am J*
18 *Med*, 118(2), 161-167 (2005).
- 19 146. Maxwell CJ, Hogan DB, Ebly EM. Serum folate levels and subsequent
20 adverse cerebrovascular outcomes in elderly persons. *Dement Geriatr Cogn*
21 *Disord*, 13(4), 225-234 (2002).

- 1 147. Ravaglia G, Forti P, Maioli F *et al.* Conversion of mild cognitive impairment to
2 dementia: predictive role of mild cognitive impairment subtypes and vascular
3 risk factors. *Dement Geriatr Cogn Disord*, 21(1), 51-58 (2006).
- 4 148. McCaddon A, Hudson P, Davies G, Hughes A, Williams JH, Wilkinson C.
5 Homocysteine and cognitive decline in healthy elderly. *Dement Geriatr Cogn*
6 *Disord*, 12(5), 309-313 (2001).
- 7 149. Seshadri S, Beiser A, Selhub J *et al.* Plasma homocysteine as a risk factor for
8 dementia and Alzheimer's disease. *N.Engl.J.Med.*, 346(7), 476-483 (2002).
- 9 150. Clarke R, Birks J, Nexø E *et al.* Low vitamin B-12 status and risk of cognitive
10 decline in older adults. *Am J Clin Nutr*, 86(5), 1384-1391 (2007).
- 11 151. Hin H, Clarke R, Sherliker P *et al.* Clinical relevance of low serum vitamin B12
12 concentrations in older people: the Banbury B12 study. *Age Ageing*, 35(4),
13 416-422 (2006).
- 14 152. Miller JW, Green R, Ramos MI *et al.* Homocysteine and cognitive function in
15 the Sacramento Area Latino Study on Aging. *Am J Clin Nutr*, 78(3), 441-447
16 (2003).
- 17 153. Haan MN, Miller JW, Aiello AE *et al.* Homocysteine, B vitamins, and the
18 incidence of dementia and cognitive impairment: results from the Sacramento
19 Area Latino Study on Aging. *Am J Clin Nutr*, 85(2), 511-517 (2007).
- 20 154. Blasko I, Jellinger K, Kemmler G *et al.* Conversion from cognitive health to
21 mild cognitive impairment and Alzheimer's disease: prediction by plasma

- 1 amyloid beta 42, medial temporal lobe atrophy and homocysteine. *Neurobiol*
2 *Aging*, 29(1), 1-11 (2008).
- 3 155. Dufouil C, Alperovitch A, Ducros V, Tzourio C. Homocysteine, white matter
4 hyperintensities, and cognition in healthy elderly people. *Ann Neurol*, 53(2),
5 214-221 (2003).
- 6 156. Luchsinger JA, Tang MX, Miller J, Green R, Mayeux R. Relation of higher
7 folate intake to lower risk of Alzheimer disease in the elderly. *Arch Neurol*,
8 64(1), 86-92 (2007).
- 9 157. Durga J, van Boxtel MP, Schouten EG *et al.* Effect of 3-year folic acid
10 supplementation on cognitive function in older adults in the FACIT trial: a
11 randomised, double blind, controlled trial. *Lancet*, 369(9557), 208-216 (2007).
- 12 158. Malouf R, Grimley Evans J. Folic acid with or without vitamin B12 for the
13 prevention and treatment of healthy elderly and demented people. *Cochrane*
14 *Database Syst Rev*, (4), CD004514 (2008).
- 15 159. Elias MF, Sullivan LM, D'Agostino RB *et al.* Homocysteine and cognitive
16 performance in the Framingham offspring study: age is important.
17 *Am.J.Epidemiol.*, 162(7), 644-653 (2005).
- 18 160. Lehmann M, Regland B, Blennow K, Gottfries CG. Vitamin B12-B6-folate
19 treatment improves blood-brain barrier function in patients with
20 hyperhomocysteinaemia and mild cognitive impairment. *Dement Geriatr Cogn*
21 *Disord*, 16(3), 145-150 (2003).

- 1 161. VanItallie TB. Subsyndromal depression in the elderly: underdiagnosed and
2 undertreated. *Metabolism*, 54(5 Suppl 1), 39-44 (2005).
- 3 162. Bottiglieri T, Laundy M, Crellin R, Toone BK, Carney MW, Reynolds EH.
4 Homocysteine, folate, methylation, and monoamine metabolism in
5 depression. *J Neurol Neurosurg Psychiatry*, 69(2), 228-232 (2000).
- 6 163. Carney MW, Chary TK, Laundy M *et al.* Red cell folate concentrations in
7 psychiatric patients. *J Affect Disord*, 19(3), 207-213 (1990).
- 8 164. Shorvon SD, Carney MW, Chanarin I, Reynolds EH. The neuropsychiatry of
9 megaloblastic anaemia. *Br Med J*, 281(6247), 1036-1038 (1980).
- 10 165. Surtees R, Heales S, Bowron A. Association of cerebrospinal fluid deficiency
11 of 5-methyltetrahydrofolate, but not S-adenosylmethionine, with reduced
12 concentrations of the acid metabolites of 5-hydroxytryptamine and dopamine.
13 *Clin Sci (Lond)*, 86(6), 697-702 (1994).
- 14 166. Bottiglieri T, Hyland K, Laundy M *et al.* Folate deficiency, biopterin and
15 monoamine metabolism in depression. *Psychol Med*, 22(4), 871-876 (1992).
- 16 167. Wesson VA, Levitt AJ, Joffe RT. Change in folate status with antidepressant
17 treatment. *Psychiatry Res*, 53(3), 313-322 (1994).
- 18 168. Godfrey PS, Toone BK, Carney MW *et al.* Enhancement of recovery from
19 psychiatric illness by methylfolate. *Lancet*, 336(8712), 392-395 (1990).
- 20 169. Rosche J, Uhlmann C, Froscher W. Low serum folate levels as a risk factor
21 for depressive mood in patients with chronic epilepsy. *J Neuropsychiatry Clin*
22 *Neurosci*, 15(1), 64-66 (2003).

- 1 170. Alpert JE, Fava M. Nutrition and depression: the role of folate. *Nutr Rev*,
2 55(5), 145-149 (1997).
- 3 171. Papakostas GI, Petersen T, Lebowitz BD *et al*. The relationship between
4 serum folate, vitamin B12, and homocysteine levels in major depressive
5 disorder and the timing of improvement with fluoxetine. *Int J*
6 *Neuropsychopharmacol*, 8(4), 523-528 (2005).
- 7 172. Papakostas GI, Petersen T, Mischoulon D *et al*. Serum folate, vitamin B12,
8 and homocysteine in major depressive disorder, Part 2: predictors of relapse
9 during the continuation phase of pharmacotherapy. *J Clin Psychiatry*, 65(8),
10 1096-1098 (2004).
- 11 173. Coppen A, Bailey J. Enhancement of the antidepressant action of fluoxetine
12 by folic acid: a randomised, placebo controlled trial. *J Affect Disord*, 60(2),
13 121-130 (2000).
- 14 174. Ramos MI, Allen LH, Haan MN, Green R, Miller JW. Plasma folate
15 concentrations are associated with depressive symptoms in elderly Latina
16 women despite folic acid fortification. *Am J Clin Nutr*, 80(4), 1024-1028
17 (2004).
- 18 175. Kendrick T, Dunn N, Robinson S *et al*. A longitudinal study of blood folate
19 levels and depressive symptoms among young women in the Southampton
20 Women's Survey. *J Epidemiol Community Health*, 62(11), 966-972 (2008).
- 21 176. Gilbody S, Lewis S, Lightfoot T. Methylene tetrahydrofolate reductase
22 (MTHFR) genetic polymorphisms and psychiatric disorders: a HuGE review.
23 *Am J Epidemiol*, 165(1), 1-13 (2007).

- 1 177. Alpert M, Silva RR, Pouget ER. Prediction of treatment response in geriatric
2 depression from baseline folate level: interaction with an SSRI or a tricyclic
3 antidepressant. *J Clin Psychopharmacol*, 23(3), 309-313 (2003).
- 4 178. Reynolds EH, Carney MW, Toone BK. Methylation and mood. *Lancet*,
5 2(8396), 196-198 (1984).
- 6 179. Otero-Losada ME, Rubio MC. Acute changes in 5-HT metabolism after S-
7 adenosyl-L-methionine administration. *Gen Pharmacol*, 20(4), 403-406
8 (1989).
- 9 180. Losada ME, Rubio MC. Acute effects of S-adenosyl-L-methionine on
10 catecholaminergic central function. *Eur J Pharmacol*, 163(2-3), 353-356
11 (1989).
- 12 181. Bottiglieri T, Godfrey P, Flynn T, Carney MW, Toone BK, Reynolds EH.
13 Cerebrospinal fluid S-adenosylmethionine in depression and dementia: effects
14 of treatment with parenteral and oral S-adenosylmethionine. *J Neurol*
15 *Neurosurg Psychiatry*, 53(12), 1096-1098 (1990).
- 16 182. Bressa GM. S-adenosyl-L-methionine (SAME) as antidepressant: meta-
17 analysis of clinical studies. *Acta Neurol Scand Suppl*, 154, 7-14 (1994).
- 18 183. Miller DB, O'Callaghan JP. Depression, cytokines, and glial function.
19 *Metabolism*, 54(5 Suppl 1), 33-38 (2005).
- 20 184. Passeri M, Cucinotta D, Abate G *et al.* Oral 5'-methyltetrahydrofolic acid in
21 senile organic mental disorders with depression: results of a double-blind
22 multicenter study. *Aging (Milano)*, 5(1), 63-71 (1993).

- 1 185. Fava M, Davidson KG. Definition and epidemiology of treatment-resistant
2 depression. *Psychiatr Clin North Am*, 19(2), 179-200 (1996).
- 3 186. Chen H, Zhang SM, Schwarzschild MA *et al*. Folate intake and risk of
4 Parkinson's disease. *Am J Epidemiol*, 160(4), 368-375 (2004).
- 5 187. de Lau LM, Koudstaal PJ, Witteman JC, Hofman A, Breteler MM. Dietary
6 folate, vitamin B12, and vitamin B6 and the risk of Parkinson disease.
7 *Neurology*, 67(2), 315-318 (2006).
- 8 188. Rogers JD, Sanchez-Saffon A, Frol AB, Diaz-Arrastia R. Elevated plasma
9 homocysteine levels in patients treated with levodopa: association with
10 vascular disease. *Arch Neurol*, 60(1), 59-64 (2003).
- 11 189. O'Suilleabhain PE, Sung V, Hernandez C *et al*. Elevated plasma
12 homocysteine level in patients with Parkinson disease: motor, affective, and
13 cognitive associations. *Arch Neurol*, 61(6), 865-868 (2004).
- 14 190. Muller T, Woitalla D, Hauptmann B, Fowler B, Kuhn W. Decrease of
15 methionine and S-adenosylmethionine and increase of homocysteine in
16 treated patients with Parkinson's disease. *Neurosci Lett*, 308(1), 54-56 (2001).
- 17 191. Muller T, Werne B, Fowler B, Kuhn W. Nigral endothelial dysfunction,
18 homocysteine, and Parkinson's disease. *Lancet*, 354(9173), 126-127 (1999).
- 19 192. Cheng H, Gomes-Trolin C, Aquilonius SM *et al*. Levels of L-methionine S-
20 adenosyltransferase activity in erythrocytes and concentrations of S-
21 adenosylmethionine and S-adenosylhomocysteine in whole blood of patients
22 with Parkinson's disease. *Exp Neurol*, 145(2 Pt 1), 580-585 (1997).

- 1 193. Zhao WQ, Latinwo L, Liu XX, Lee ES, Lamango N, Charlton CG. L-dopa
2 upregulates the expression and activities of methionine adenosyl transferase
3 and catechol-O-methyltransferase. *Exp Neurol*, 171(1), 127-138 (2001).
- 4 194. Muller T, Woitalla D, Fowler B, Kuhn W. 3-OMD and homocysteine plasma
5 levels in parkinsonian patients. *J Neural Transm*, 109(2), 175-179 (2002).
- 6 195. Lamberti P, Zoccolella S, Iliceto G *et al.* Effects of levodopa and COMT
7 inhibitors on plasma homocysteine in Parkinson's disease patients. *Mov*
8 *Disord*, 20(1), 69-72 (2005).
- 9 196. Irizarry MC, Gurol ME, Raju S *et al.* Association of homocysteine with plasma
10 amyloid beta protein in aging and neurodegenerative disease. *Neurology*,
11 65(9), 1402-1408 (2005).
- 12 197. Korten A, Lodder J, Vreeling F, Boreas A, van Raak L, Kessels F. Stroke and
13 idiopathic Parkinson's disease: does a shortage of dopamine offer protection
14 against stroke? *Mov Disord*, 16(1), 119-123 (2001).
- 15 198. McCann SJ, LeCouteur DG, Green AC *et al.* The epidemiology of Parkinson's
16 disease in an Australian population. *Neuroepidemiology*, 17(6), 310-317
17 (1998).
- 18 199. Emre M. Dementia associated with Parkinson's disease. *Lancet Neurol*, 2(4),
19 229-237 (2003).
- 20 200. Ozer F, Meral H, Hanoglu L *et al.* Plasma homocysteine levels in patients
21 treated with levodopa: motor and cognitive associations. *Neurol Res*, 28(8),
22 853-858 (2006).

- 1 201. Zoccolella S, Lamberti P, Iliceto G *et al.* Plasma homocysteine levels in L-
2 dopa-treated Parkinson's disease patients with cognitive dysfunctions. *Clin*
3 *Chem Lab Med*, 43(10), 1107-1110 (2005).
- 4 202. Applebaum J, Shimon H, Sela BA, Belmaker RH, Levine J. Homocysteine
5 levels in newly admitted schizophrenic patients. *J Psychiatr Res*, 38(4), 413-
6 416 (2004).
- 7 203. Goff DC, Bottiglieri T, Arning E *et al.* Folate, homocysteine, and negative
8 symptoms in schizophrenia. *Am J Psychiatry*, 161(9), 1705-1708 (2004).
- 9 204. Shi J, Gershon ES, Liu C. Genetic associations with schizophrenia: Meta-
10 analyses of 12 candidate genes. *Schizophr Res*, 104(1-3), 96-107 (2008).
- 11 205. Miodownik C, Lerner V, Vishne T, Sela BA, Levine J. High-dose vitamin B6
12 decreases homocysteine serum levels in patients with schizophrenia and
13 schizoaffective disorders: a preliminary study. *Clin Neuropharmacol*, 30(1),
14 13-17 (2007).
- 15 206. Levine J, Stahl Z, Sela BA *et al.* Homocysteine-reducing strategies improve
16 symptoms in chronic schizophrenic patients with hyperhomocysteinemia. *Biol*
17 *Psychiatry*, 60(3), 265-269 (2006).
- 18 207. Ozbek Z, Kucukali CI, Ozkok E *et al.* Effect of the methylenetetrahydrofolate
19 reductase gene polymorphisms on homocysteine, folate and vitamin B12 in
20 patients with bipolar disorder and relatives. *Prog Neuropsychopharmacol Biol*
21 *Psychiatry*, 32(5), 1331-1337 (2008).

- 1 208. Reif A, Pfuhlmann B, Lesch KP. Homocysteinemia as well as
2 methylenetetrahydrofolate reductase polymorphism are associated with
3 affective psychoses. *Prog Neuropsychopharmacol Biol Psychiatry*, 29(7),
4 1162-1168 (2005).
- 5 209. Dittmann S, Seemuller F, Grunze HC *et al.* The impact of homocysteine levels
6 on cognition in euthymic bipolar patients: a cross-sectional study. *J Clin*
7 *Psychiatry*, 69(6), 899-906 (2008).
- 8 210. Dittmann S, Seemuller F, Schwarz MJ *et al.* Association of cognitive deficits
9 with elevated homocysteine levels in euthymic bipolar patients and its impact
10 on psychosocial functioning: preliminary results. *Bipolar Disord*, 9(1-2), 63-70
11 (2007).
- 12 211. Osher Y, Sela BA, Levine J, Belmaker RH. Elevated homocysteine levels in
13 euthymic bipolar disorder patients showing functional deterioration. *Bipolar*
14 *Disord*, 6(1), 82-86 (2004).
- 15 212. Miller A, Korem M, Almog R, Galboiz Y. Vitamin B12, demyelination,
16 remyelination and repair in multiple sclerosis. *J Neurol Sci*, 233(1-2), 93-97
17 (2005).
- 18 213. Vrethem M, Mattsson E, Hebelka H *et al.* Increased plasma homocysteine
19 levels without signs of vitamin B12 deficiency in patients with multiple
20 sclerosis assessed by blood and cerebrospinal fluid homocysteine and
21 methylmalonic acid. *Mult Scler*, 9(3), 239-245 (2003).

- 1 214. Ramsaransing GS, Fokkema MR, Teelken A, Arutjunyan AV, Koch M, De
2 Keyser J. Plasma homocysteine levels in multiple sclerosis. *J Neurol*
3 *Neurosurg Psychiatry*, 77(2), 189-192 (2006).
- 4 215. Reynolds EH, Bottiglieri T, Laundry M, Crellin RF, Kirker SG. Vitamin B12
5 metabolism in multiple sclerosis. *Arch Neurol*, 49(6), 649-652 (1992).
- 6 216. Goodkin DE, Jacobsen DW, Galvez N, Daughtry M, Secic M, Green R. Serum
7 cobalamin deficiency is uncommon in multiple sclerosis. *Arch Neurol*, 51(11),
8 1110-1114 (1994).
- 9 217. Schroecksnadel K, Frick B, Wirleitner B, Winkler C, Schennach H, Fuchs D.
10 Moderate hyperhomocysteinemia and immune activation. *Curr Pharm*
11 *Biotechnol*, 5(1), 107-118 (2004).
- 12 218. Lawson BR, Manenkova Y, Ahamed J *et al.* Inhibition of transmethylation
13 down-regulates CD4 T cell activation and curtails development of
14 autoimmunity in a model system. *J Immunol*, 178(8), 5366-5374 (2007).
- 15 219. Wade DT, Young CA, Chaudhuri KR, Davidson DL. A randomised placebo
16 controlled exploratory study of vitamin B-12, lofepramine, and L-phenylalanine
17 (the "Cari Loder regime") in the treatment of multiple sclerosis. *J Neurol*
18 *Neurosurg Psychiatry*, 73(3), 246-249 (2002).
- 19 220. van Rensburg SJ, Kotze MJ, Hon D *et al.* Iron and the folate-vitamin B12-
20 methylation pathway in multiple sclerosis. *Metab Brain Dis*, 21(2-3), 121-137
21 (2006).

- 1 221. Grandes P, Kq K, Morino P, Cuenod M, Streit P. Homocysteate, an Excitatory
2 Transmitter Candidate Localized in Glia. *Eur J Neurosci*, 3(12), 1370-1373
3 (1991).
- 4 222. Neubauer C. Mental deterioration in epilepsy due to folate deficiency. *Br Med*
5 *J*, 2(5712), 759-761 (1970).
- 6 223. Schwaninger M, Ringleb P, Winter R *et al*. Elevated plasma concentrations of
7 homocysteine in antiepileptic drug treatment. *Epilepsia*, 40(3), 345-350
8 (1999).
- 9 224. Sener U, Zorlu Y, Karaguzel O, Ozdamar O, Coker I, Topbas M. Effects of
10 common anti-epileptic drug monotherapy on serum levels of homocysteine,
11 vitamin B12, folic acid and vitamin B6. *Seizure*, 15(2), 79-85 (2006).
- 12 225. Billings RE. Decreased hepatic 5, 10-methylenetetrahydrofolate reductase
13 activity in mice after chronic phenytoin treatment. *Mol Pharmacol*, 25(3), 459-
14 466 (1984).
- 15 226. Morrell MJ. Guidelines for the care of women with epilepsy. *Neurology*, 51(5
16 Suppl 4), S21-27 (1998).
- 17 227. Gidal BE, Tamura T, Hammer A, Vuong A. Blood homocysteine, folate and
18 vitamin B-12 concentrations in patients with epilepsy receiving lamotrigine or
19 sodium valproate for initial monotherapy. *Epilepsy Res*, 64(3), 161-166
20 (2005).

- 1 228. Karabiber H, Sonmezgoz E, Ozerol E, Yakinci C, Otlu B, Yologlu S. Effects of
2 valproate and carbamazepine on serum levels of homocysteine, vitamin B12,
3 and folic acid. *Brain Dev*, 25(2), 113-115 (2003).
- 4 229. Crawford P. Best practice guidelines for the management of women with
5 epilepsy. *Epilepsia*, 46 Suppl 9, 117-124 (2005).
- 6 230. Wilson RD, Davies G, Desilets V *et al.* The use of folic acid for the prevention
7 of neural tube defects and other congenital anomalies. *J Obstet Gynaecol*
8 *Can*, 25(11), 959-973 (2003).
- 9 231. Monji A, Yanagimoto K, Maekawa T, Sumida Y, Yamazaki K, Kojima K.
10 Plasma folate and homocysteine levels may be related to interictal
11 "schizophrenia-like" psychosis in patients with epilepsy. *J Clin*
12 *Psychopharmacol*, 25(1), 3-5 (2005).
- 13 232. Edeh J, Toone BK. Antiepileptic therapy, folate deficiency, and psychiatric
14 morbidity: a general practice survey. *Epilepsia*, 26(5), 434-440 (1985).
- 15 233. Ali A, Pillai KK, Pal SN. Effects of folic acid and lamotrigine therapy in some
16 rodent models of epilepsy and behaviour. *J Pharm Pharmacol*, 55(3), 387-391
17 (2003).
- 18 234. Huemer M, Ausserer B, Graninger G *et al.* Hyperhomocysteinemia in children
19 treated with antiepileptic drugs is normalized by folic acid supplementation.
20 *Epilepsia*, 46(10), 1677-1683 (2005).

- 1 235. Apeland T, Mansoor MA, Pentieva K, McNulty H, Seljeflot I, Strandjord RE.
2 The effect of B-vitamins on hyperhomocysteinemia in patients on antiepileptic
3 drugs. *Epilepsy Res*, 51(3), 237-247 (2002).
- 4 236. Reynolds EH. Mental effects of anticonvulsants, and folic acid metabolism.
5 *Brain*, 91(2), 197-214 (1968).
- 6 237. Smith AD. Folic acid fortification: the good, the bad, and the puzzle of vitamin
7 B-12. *Am J Clin Nutr*, 85(1), 3-5 (2007).
- 8 238. Luebeck EG, Moolgavkar SH, Liu AY, Boynton A, Ulrich CM. Does folic acid
9 supplementation prevent or promote colorectal cancer? Results from model-
10 based predictions. *Cancer Epidemiol Biomarkers Prev*, 17(6), 1360-1367
11 (2008).
- 12 239. Smith AD, Kim YI, Refsum H. Is folic acid good for everyone? *Am J Clin Nutr*,
13 87(3), 517-533 (2008).
- 14 240. Pitkin RM. Folate and neural tube defects. *Am J Clin Nutr*, 85(1), 285S-288S
15 (2007).
- 16 241. Molloy AM, Kirke PN, Troendle JF *et al*. Maternal vitamin B12 status and risk
17 of neural tube defects in a population with high neural tube defect prevalence
18 and no folic Acid fortification. *Pediatrics*, 123(3), 917-923 (2009).
- 19 242. Butterworth CE, Jr., Tamura T. Folic acid safety and toxicity: a brief review.
20 *Am J Clin Nutr*, 50(2), 353-358 (1989).
- 21 243. Herzlich B, Herbert V. Depletion of serum holotranscobalamin II. An early sign
22 of negative vitamin B12 balance. *Lab Invest*, 58(3), 332-337 (1988).